

# Diagnostic Accuracy of Fine Needle Aspiration Cytology, Triple Test and Tru-cut Biopsy in the Detection of Breast Lesion

Nirmal Kumar Agarwal, Dharendra Nath Choudhury, Akshit Minocha

Department of General Surgery, India.

## Abstract

**Background:** The majority of breast illnesses manifest as a palpable mass. The vast majority of breast lesions are not malignant, and the vast majority of benign lesions do not progress to cancer; however, the accuracy of diagnosis can be improved by a combination of preoperative tests such as physical examination, mammography, fine-needle aspiration cytology (FNAC), and Trucut needle biopsy (TCNB) or core needle biopsy (CNB). FNAC has gained in popularity and is now the primary treatment performed after a history and clinical examination to diagnose solid and cystic breast masses. This study was designed to assess the accuracy of FNAC in comparison to Triple Test and TCNB for the diagnosis of breast masses. **Methodology:** From January 2023 through May 2023, this study was conducted in the Tezpur Medical College and Hospital. The research included all female patients over the age of 20 with clinically palpable lumps. Palpable breast lumps need a complete clinical breast examination, imaging, and tissue sample for a clear diagnosis to rule out cancer. The Ultrasonography/Mammography, FNAC and tru-cut biopsy procedures were performed concurrently in the same region, and the results were compared to final histopathology as the gold standard. **Result:** The majority of the patients in the study were in age group of > 60 years. In our study Left breast was involved in 41 (58.5%) patients and right breast in 29 (41.4%) patients. The most frequent site involved was upper outer quadrant in 38 (54.2%). The overall sensitivity of FNAC in our study was 93.75%, tru-cut biopsy is 100% and that of triple test is 100% and specificity for malignancy was 91.6%, 93.6 and 95.6% for FNAC, tru-cut biopsy and triple test respectively. **Conclusion:** All three techniques have advantages and disadvantages. Core Needle Biopsy cannot be used in place of FNAC, and it is not required to identify all breast lesions. It can be utilised as an adjuvant in circumstances when cytology is definitive and biomarker tests are required. triple test and trucut were more accurate than FNAC in our testing. Trucut biopsy was able to provide histological diagnosis, and the findings were 100% consistent with the final histopathology report.

**Keywords:** Fine needle aspiration cytology (FNAC)- Tru-cut needle biopsy (TCNB)- Triple Test- breast lump

*Asian Pac J Cancer Care*, 8 (3), 539-543

Submission Date: 06/22/2023

Acceptance Date: 08/18/2023

## Introduction

Breast cancer is one of the most investigated cancers in the world, and new developments in its management are common. In India, the prevalence of breast cancer has now reached ENDEMIC levels. Numerous disorders, most of which manifest as lumps in the breast, including benign and malignant neoplasms, inflammatory diseases, and infections, can affect the breast. Breast lumps are among the most frequent complaints in surgical OPDs, thus it's critical to distinguish between benign and malignant disorders before administering treatment. A comprehensive

examination is the key to accurate diagnosis. Twenty to thirty percent of all malignancies worldwide [1] are caused by it. Early diagnosis is of utmost importance in the medical industry because treating patients in latter stages is frequently ineffective. Lack of education and screening programmes causes the ignorant people to be unaware of the terrible disease. The goal of developing a sensitive, specific, effective, and affordable way to diagnose breast cancer was to examine the breast lumps using a variety of diagnostic techniques. examination of

### Corresponding Author:

Dr. Akshit Minocha

Department of General Surgery, India.

Email: akshitminocha1@gmail.com

the breast, mammography, Trucut biopsy (needle core biopsy), ultrasonography, thermography, FNAC, open excision biopsy is all used in diagnostic work up of a palpable breast lump [2]. This goal has long been served by FNAC, in addition to clinical evaluation and mammography. It has been shown to be quite useful in the diagnosis of breast lumps since, in addition to being affordable, it also makes the cytological testing [3] process simple and quick. The first priority investigation for patients with breast lumps is frequently the FNAC. Nevertheless, it has some drawbacks, including the inability to distinguish between invasive and in situ carcinomas, a lack of sample size, and false negative results. One of the important methods for obtaining a histopathological diagnosis today is trucut biopsy, also known as core needle biopsy. It is manageable and can be done without hospitalisation.

The triple test, first reported in 1975, involves the physical examination, mammography, and fine-needle aspiration of palpable breast tumours. When all three components are concordant, that is, all benign or all malignant, the triple test score (TTS) is helpful and reliable for assessing palpable breast masses and can substitute open surgical biopsy for diagnosis [4]. If the findings are inconsistent, the patient may be submitted to further testing and an open surgical biopsy. Clinical breast examination, breast imaging (Breast Ultrasound and/or Mammogram), and fine needle aspiration cytology are the three components of the Triple test.

A triple test was also submitted to a score system for the parameters mentioned earlier. Each criterion receives one to three points: benign (1 point), suspicious (2 points), and malignant (3 points). The total score between 3 and 9 is used to interpret the results.

1. A total score of 3-4 indicates a benign lesion.

2. A total score of 5 indicates an intermediate risk that necessitates an excisional biopsy.

3. A total score of 6 or above indicates the possibility of cancer and the need for surgical intervention.

**TRU-CUT Biopsy:** When evaluating a discrete lump in the breast, an ultrasound-guided automated Tru-cut needle biopsy may be utilised instead of fine needle aspiration cytology. The procedure's sensitivity for diagnosing substantial pathology varies from 88.7% to 97%, while its specificity ranges from 96.8% to 100% [5-7]. When utilised in conjunction with a triple assessment, the sensitivity rises to 97.9% [7].

Core biopsy (CB) usage has grown, although not necessarily for evidence-based reasons. CB and FNAC do not contradict one other. FNAC should be used in the diagnosis of benign, symptomatic lesions, and CB as an alternative diagnostic modality should be used with caution, e.g., in the diagnosis of impalpable masses, microcalcifications, or a clinically apparent malignancy where preoperative chemotherapy is planned. CB should not be utilised to compensate for poor FNAC performance. Where expert cytopathologists are available, FNAC and CB may complement each other to give a highly accurate, quick, and cost-effective method of patient triage [8].

## Materials and Methods

From January 2023 to May 2023, 70 consecutive females with palpable breast lumps were examined at the surgical out-patient department of the Tezpur Medical College and Hospital. The study included all girls older than 20 with palpable breast masses. Men, impalpable masses, recurring cancer, and age less than 20 years. All patients provided written informed consent, which was obtained. Each patient underwent FNAC and tru-cut biopsy simultaneously after a thorough history, clinical examination, and mammography in those patients (> 40 years) who were indicated. A skilled surgeon performed the tru-cut biopsy, while a histopathologist performed the FNAC. To reduce the danger of bias, the FNAC and tru-cut biopsy slides were examined by two different histopathologists. Later, all of them underwent mastectomy or excision biopsy of the lump for definitive diagnosis confirmation. Both techniques results were collected, and their sensitivity levels were computed and compared.

FNAC was performed using an aseptic approach and a disposable syringe with a 21-23 gauge needle. Under local anaesthesia, tru-cut biopsy was performed on the lesions using needles between 14 and 16 gauge.

## Results

In this study 70 females underwent FNAC, Tru-cut biopsy and Triple Assessment simultaneously. Mean age was 51.1

Left breast was involved in 41 (58.5%) patients and right breast in 29 (41.4%) patients. The most frequent site involved was upper outer quadrant in 38 (54.2%) patients central in 18 (25.7%) patients lower inner in 8 (11.4%) patients and upper inner in 6 (8.5%) respectively.

FNAC diagnosed breast cancer in 24 patients and 4 cases one of benign phyllodes, one atypical epithelial hyperplasia and two giant fibroadenoma were reported false positive. Four cases of invasive ductal carcinomas were reported false negative. Fibroadenomas were diagnosed in 26 cases, 24 confirmed on biopsy and one found out to be phylloides tumor and other epithelial hyperplasia each (false positive).

Out of 70 breast lesion 60 were diagnosed correctly on FNAC with 93.75% sensitivity and specificity for malignancy was 91.6% (Table 1).

Trucut biopsy diagnosed the carcinoma in 26 patients. Invasive ductal carcinoma was diagnosed correctly in

Table 1. Showing Number of Breast Lump Cases in Different Age Groups

Age group (in years)	Number of cases
<30	6
30-39	8
40-49	14
50-59	20
>60	22

Table 2. Showing Number of Patients as Per Different Category of Breast Lesion after FNAC

FNAC Diagnosis	Number of cases	Final Histopathology	False Positive	False Negative
Benign	20	Fibroadenoma - 11	-	4
		Phylloides Tumor - 3		
		Epithelial Hyperplasia -2		
		Ductal Cancer -4		
Fibroadenoma	26	Fibroadenoma -22	2	-
		Phylloides tumor -1		
		Epithelial hyperplasia-1		
Ductal carcinoma	3	Ductal cancer - 3		
Malignant	21	Ductal cancer - 13	4	-
		Lobular Cancer - 4		
		Atypical Epithelial Hyperplasia -1		
		Phylloides -1		
		Giant Fibroadenoma -2		

Table 3. Showing Number of Patients as Per Different Category of Breast Lesion after Tru-cut Biopsy

Tru-cut Diagnosis	Number of patients	Final Histopathology	False Positive	False Negative
Benign	8	Fibroadenoma - 4	-	-
		Phylloides Tumor - 2		
		Epithelial Hyperplasia -2		
Fibroadenoma	34	Fibroadenoma - 34	-	-
Phylloides	2	Phylloides -2	-	-
Ductal Cancer	16	Ductal Cancer - 16	-	-
Malignant	10	Ductal cancer - 4	3	-
		Lobular Cancer - 3		
		Atypical Epithelial Hyperplasia -1		
		Phylloides -1		
		Giant Fibroadenoma -1		

16 cases. 10 cases which were only reported malignant on tru-cut came out to be invasive ductal carcinoma in 4 cases, lobular carcinoma in 3 cases and each case of benign phylloides tumor, atypical epithelial hyperplasia and giant fibroadenoma were false positive. 34 cases of fibroadenoma diagnosed on tru-cut were confirmed accurate on biopsy. Sensitivity calculated is 100%. with sixty seven out of seventy were reported accurately with no false negative result and specificity for malignancy is 93.6% (Table 2).

Triple Test diagnosed the carcinoma in 26 patients. Invasive ductal carcinoma was diagnosed correctly in 17 cases. 9 cases which were only reported malignant on tru-cut came out to be invasive ductal carcinoma in 4 cases, lobular carcinoma in 2 cases and each case of benign phylloides tumor and giant fibroadenoma were false positive. 34 cases of fibroadenoma diagnosed on Triple Test confirmed accurate on biopsy. Sensitivity calculated is 100%. with sixty eight out of seventy were reported accurately with no false negative result and specificity for malignancy is 95.6% (Table 3).

The overall sensitivity of FNAC in our study was 93.75%, tru-cut biopsy is 100% and that of triple test is 100% and specificity for malignancy was 91.6%, 93.6 and 95.6% for FNAC, tru-cut biopsy and triple test

respectively.

## Discussion

In our study, FNAC and trucut biopsy were performed on the same lesion and the operator dependence has been standardized. People used to undergo excision biopsy of fibroadenoma to remove the tumor and establish diagnosis. Now tru-cut biopsy is widely used and lesion is left undisturbed in the breast if the diagnosis is fibroadenoma. Fibroadenoma was the most common histopathological diagnosis and comprised 54.2% of the total cases. In our study, 38 cases of fibroadenomas were reported on histopathology, of them 34 were correctly diagnosed on Tru-cut biopsy reflecting the high accuracy of trucut biopsy (Table 3). In 3 (4.2%), 16 (22.8%) and 17 (24.2%) of these were detected on FNAC, Trucut samples and Triple test accurately detected invasive ductal cancer. The remaining 13 (18.5%), 4 (6.1%) and 4 cases in FNAC Trucut biopsies and triple test, respectively, were reported as malignant (Tables 2, 3 and 4). Four incidences of invasive ductal carcinoma were identified on histology out of 20 benign lesions on FNAC (Table 2). respectively. "Phylloides tumours can spread to other parts of the body if they are malignant and have a 20% to 40% local recurrence rate [9].

Table 4. Showing Number of Patients as Per Different Category of Breast Lesion after Triple Test

Triple Test Diagnosis	Number of patients	Final Histopathology	False Positive	False Negative
Benign	8	Fibroadenoma - 4 Phylloides Tumor - 2 Epithelial Hyperplasia -2	-	-
Fibroadenoma	34	Fibroadenoma - 34	-	-
Phylloides	2	Phylloides -2	-	-
Ductal Cancer	17	Ductal Cancer - 17	-	-
Malignant	19	Ductal cancer - 4 Lobular Cancer - 3 Phyllodes -1 Giant Fibroadenoma -1	2	-

In our investigation, two out of five phylloides tumours had a correct Tru-cut biopsy diagnosis. The remaining three were described as benign (two) and malignant (one). FNAC identified three of the phylloides tumours as benign lesions, one as a fibroadenoma, and one as malignant after failing to diagnose any of the tumours. Overall FNAC sensitivity in our study was 93.75%, and trucut biopsy sensitivity was 100%. In a study that was equivalent to ours, the sensitivity of FNAC and Trucut was found to be 88% and 96%, respectively [10]. Our study's results are comparable to those from another study [11] that indicated the FNAC and trucut biopsy's sensitivity as 81.4% and 91.5%, respectively. Our study's 93.75% FNAC sensitivity was equivalent to those of prior research [12-14].” The 100% sensitivity of the trucut biopsy in our study is comparable to the 96.7% and 100% sensitivities of the trucut biopsy in studies by Loffe et al [15] and Memon et al [16]. In our investigation, the FNAC and trucut specificities for malignancy were 91.6% and 93.6%, respectively, comparable to the FNAC (100%) and trucut biopsy (100%) in one study [4,17].

In conclusion, all three techniques have advantages and disadvantages. Core Needle Biopsy cannot be used in place of FNAC, and it is not required to identify all breast lesions. It can be utilised as an adjuvant in circumstances when cytology is definitive and biomarker tests are required. Triple test and trucut were more accurate than FNAC in our testing. Trucut biopsy was able to provide histological diagnosis, and the findings were 100% consistent with the final histopathology report.

## References

- Masood S, Frykberg ER, McLellan GL, Scalapino MC, Mitchum DG, Bullard JB. Prospective evaluation of radiologically directed fine-needle aspiration biopsy of nonpalpable breast lesions. *Cancer*. 1990 Oct 01;66(7):1480-1487. [https://doi.org/10.1002/1097-0142\(19901001\)66:7<1480::aid-cncr2820660708>3.0.co;2-o](https://doi.org/10.1002/1097-0142(19901001)66:7<1480::aid-cncr2820660708>3.0.co;2-o)
- Muhammad E, Ahmed A, Osman S. Validity of Fine Needle Aspiration Cytology (FNAC) in Diagnosis of Breast Lumps in Upper Egypt. *Egyptian Journal of Medical Laboratory Sciences*. 2012 01 01;21:69-81.
- Patel JJ, Gartell PC, Smallwood JA, Herbert A, Royle G, Buchanan R, Taylor I. Fine needle aspiration cytology of breast masses: an evaluation of its accuracy and reasons for diagnostic failure. *Annals of the Royal College of Surgeons of England*. 1987 07;69(4):156-159.
- Bdour M, Hourani S, Mefleh W, Shabatat A, Karadsheh S, Nawaiseh O, et al. Comparison between fine needle aspiration cytology and tru-cut biopsy in the diagnosis of breast cancer. *J of Surg Pak*. 2008;13(1):19-21.
- Gukas ID, Nwana EJ, Ihezue CH, Momoh JT, Obekpa PO. Tru-cut biopsy of palpable breast lesions: a practical option for pre-operative diagnosis in developing countries. *The Central African Journal of Medicine*. 2000 05;46(5):127-130. <https://doi.org/10.4314/cajmv.v46i5.8536>
- Woodcock NP, Graves I, Morgan DR, MacFie J. Ultrasound-guided Tri-cut biopsy of the breast. *Annals of the Royal College of Surgeons of England*. 1998 07;80(4):253-256.
- Nauvi SR, Ahmed TM, Naqvi SS, Jan B. Comparison of diagnostic accuracy of core biopsy for breast lesions with FNAC. *Pakistan Armed Forces Medical Journal*. 2010;01:1-3.
- Bak M, Szabó E, Mándoky L. [The “gray zone” in fine needle aspiration cytology of the breast]. *Magyar Sebeszet*. 2005 02;58(1):3-7.
- Rosen PP, Oberman HA. Fibroepithelial neoplasms. In: *Autor. Atlas of tumor pathology -tumors of the mammary gland*. 3<sup>rd</sup> ed. Washington: Armed Forces Institute of Pathology. 1993;101-14.
- Jalali U, Rasul S, Khan A. Comparison of trucut biopsy with fine needle aspiration cytology in diagnosis of palpable breast lesions. *JSurgPak*. 2004;9:27-30.
- Ghorori RMH, Ewais MK. Fine needle aspiration cytology vs core needle biopsy in diagnosis of benign breast diseases. *AAMJ*. 2004;2:127-42.
- Horgan PG, Waldron D, Mooney E, O'Brien D, McGuire M, Given HF. The role of aspiration cytologic examination in the diagnosis of carcinoma of the breast. *Surgery, Gynecology & Obstetrics*. 1991 04;172(4):290-292.
- Park IA, Ham EK. Fine needle aspiration cytology of palpable breast lesions. Histologic subtype in false negative cases. *Acta Cytologica*. 1997;41(4):1131-1138. <https://doi.org/10.1159/000332799>
- Aziz M, Ahmad N, Zahid J, Faizullah N, Aziz M. Comparison of FNAC and open biopsy in palpable breast lumps. *Journal of the College of Physicians and Surgeons--Pakistan: JCPSP*. 2004 Nov;14(11):654-656.
- Ioffe OB, Berg WA, Silverberg SG, Kumar D. Mammographic-histopathologic correlation of large-core needle biopsies of the breast. *Modern Pathology: An Official Journal of the United States and Canadian Academy of Pathology, Inc*. 1998 08;11(8):721-727.
- Memon F, Qureshi S, Zubair M, Baloch S, Quraishy MS. Sensitivity of fine needle aspiration vs core biopsy in the

diagnosis of palpable and clinically suspicious breast lesions.  
PJS . 2009;25:214-7.

17. Kumar SS, Grace RD, Babu KD, Bhaskar M. A comparative study of fine needle aspiration cytology, trucut biopsy and histopathological examination in breast lump. IOSR-JDMS . 2015;14:42-5.



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.