

# Intracranial Hemangiopericytoma- A Case Report

Mayur Khandelwal, Priya Tawri, Shankar Lal Jakhar

Department of Radiation Oncology, Acharya Tulsi Regional Cancer Institute, Sardar Patel Medical College, Bikaner (Rajasthan), India.

## Abstract

Intracranial hemangiopericytomas are rare vascular tumours, account for less than 1% of all primary tumours of the central nervous system. This tumour usually found outside of CNS i.e., lower extremities and retroperitoneum. Treatment is surgery followed by radiotherapy to decrease chances of recurrences. This is a case report of a young female diagnosed with intracranial hemangiopericytoma in 2021 at our hospital-Acharya Tulsi Regional Cancer Institute, Bikaner.

**Keywords:** Intracranial hemangiopericytoma- Radiotherapy- Recurrences

*Asian Pac J Cancer Care*, 9 (1), 203-205

Submission Date: 10/13/2023    Acceptance Date: 12/10/2023

## Introduction

Hemangiopericytomas were described by stout as malignant tumours which arises from blood vessels and he described these as sarcoma [1]. The cell of origin of these HPCs are pericytes which interacts with endothelial cells and critical for blood vessel formation and maintenance [2]. HPCs are rare tumours occurs mostly outside CNS only (in lower extremities and retroperitoneum) but sometimes also seen in CNS [3]. A case series done by Guthrie et al. [4] showed that most patients presenting with meningeal HPCs are adults, usually 30-45 year of age. Hemangiopericytoma is more commonly located in supratentorial region. Slight male predominance is seen in this tumour (1.4:1). Presenting Symptoms depends on site of mass and when it arises from CNS then common symptoms are headache, seizure, visual dysfunction, and motor weakness. HPCs are significantly less common than meningioma and their clinical behaviour is more aggressive than that of benign meningioma. Moreover, they have a strong tendency of local recurrence and extracranial metastasis

### Case Report

•A 35-year-old female was admitted to our hospital (ATRCTRI Bikaner) for a sudden seizure episode for the first time. She had no previous history of seizure or family history of the disease. She had history of weakness and numbness in her left lower limb for last 3 years. The

patient had no history of trauma or fever. The neurological examination was done and we found that power in right limb was decreased compare to left and atrophy was seen in limbs.

•CECT brain was done and it showed approx. 5x3 cm Dural based sphenoid wing mass lesion seen. surgery was done to remove the mass and the specimen was sent to our pathology laboratory for histopathological examination.

• Histological examination showed proliferation of a staghorn vascular pattern of spindle cells with moderate amount of eosinophilic cytoplasm but necrosis was not seen in prepared specimen (Figure 1 and 2).

### Workup

•Patient underwent routine investigations CBC, RFT, USG and CT which showed a lesion in right side causing contralateral midline deviation. After diagnosis, patient underwent tumor embolization and complete surgical excision in February, 2021.

• An Immunohistochemistry of the specimen came positive for STAT6, CD99, and CD34.

• He was started on Adjuvant Radiotherapy 60 Gy/30# in March 2021.

• He remained in clinical follow up with our department, and is clinically stable and showed good response.

## Corresponding Author:

Dr. Mayur Khandelwal

Department of Radiation Oncology, Acharya Tulsi Regional Cancer Institute, Sardar Patel Medical College, Bikaner (Rajasthan), India.

Email: drmayur1@yahoo.com




Figure 1. Contouring at TPS on Right Side of Skull (post op volume) with Staghorn Pattern Seen in Pathology Slide.



Figure 2. Contouring is Shown on TPS with Outer Blue Line Shows PTV and Inner Yellow Line Denotes CTV.

## Discussion

HPC is a rare neoplasm, arises from pericytes, a supportive cell of the vasculature. Usually, HPCs arise in lower extremities or retroperitoneum, but rarely, it can occur within the CNS also. HPCs were previously known as angioblastic meningiomas, as the tumors are very vascular and arose from the meninges [5]. Usual presentation of patient in adulthood, with median occurrence of HPC is in the fifth decade of life. In our case the patient age is 35 year. Initially patient present with focal neurological signs and symptoms and that helps in diagnosis of disease. Recurrence rate of tumor is about 60 to 76 %. Very rarely it can also metastasize to other organ like bone, liver lung and peritoneum with 5-year metastasis rate at 33% [4]. Treatment of meningeal HPCs is surgery like meningiomas but because of extensive vasculature most of the time only surgery is not possible and pre-embolization of the tumor before resection can prevent blood loss [6]. Post op radiotherapy can be given to prevent recurrence, but a study done by Dufour et al. [7] showed that radiotherapy did not change the rate of metastases. In our case patient underwent craniotomy and then after 25 days of surgery patient came for post operative radiotherapy. Patient was treated with 60 Gy in 30 fractions and after 6 months of follow up patient was asymptomatic. Use of chemotherapy like Doxorubicin-based regimens have been used, but have yet to establish

a clear role in the treatment of HPC [8]

Therefore, we conclude that primary treatment for HPCs are surgery followed by radiotherapy. Chemotherapy has limited role here. For recurrent HPCs Dasatinib can be used because Dasatinib can inhibit PDGFR and it has been shown that overexpression of PDGFR is seen in HPCs [9].

### Patient consent declaration

We obtained patient consent for images and other clinical information to reporting in the article. The patient understands that his names and initials will not be published.

## References

1. Koch M, Nielsen GP, Yoon SS. Malignant tumors of blood vessels: angiosarcomas, hemangioendotheliomas, and hemangiopericytomas. *Journal of Surgical Oncology*. 2008 03 15;97(4):321-329. <https://doi.org/10.1002/jso.20973>
2. Bergers G, Song S. The role of pericytes in blood-vessel formation and maintenance. *Neuro-Oncology*. 2005 Oct;7(4):452-464. <https://doi.org/10.1215/S1152851705000232>
3. Espat NJ, Lewis JJ, Leung D, Woodruff JM, Antonescu CR, Shia J, Brennan MF. Conventional hemangiopericytoma: modern analysis of outcome. *Cancer*. 2002 Oct 15;95(8):1746-1751. <https://doi.org/10.1002/cncr.10867>
4. Guthrie BL, Ebersold MJ, Scheithauer BW, Shaw EG. Meningeal hemangiopericytoma: histopathological features, treatment..
5. Jaaskelainen J, Louis DN, Paulus W. World Health Organization classification of tumours: pathology and genetics of tumours of the nervous system; in Kleihues P, Cavenee WK (eds): *Haemangiopericytoma*. Lyon. 2000;:190-2.
6. Fountas KN, Kapsalaki E, Kassam M, Feltes CH, Dimopoulos VG, Robinson JS, Smith JR. Management of intracranial meningeal hemangiopericytomas: outcome and experience. *Neurosurgical Review*. 2006 04;29(2):145-153. <https://doi.org/10.1007/s10143-005-0001-9>
7. Dufour H, Métellus P, Fuentes S, Murracchiole X, Régis J, Figarella-Branger D, Grisoli F. Meningeal hemangiopericytoma: a retrospective study of 21 patients with special review of postoperative external radiotherapy. *Neurosurgery*. 2001 04;48(4):756-762; discussion 762-763. <https://doi.org/10.1097/00006123-200104000-00011>
8. Galanis E, Buckner JC, Scheithauer BW, Kimmel DW, Schomberg PJ, Piepgras DG. Management of recurrent

meningeal hemangiopericytoma. *Cancer*. 1998 05 15;82(10):1915-1920.

9. Dietzmann K, Bossanyi P, Warich-Kirches M, Kirches E, Synowitz HJ, Firsching R. Immunohistochemical detection of vascular growth factors in angiomatous and atypical meningiomas, as well as hemangiopericytomas. *Pathology, Research and Practice*. 1997;193(7):503-510. [https://doi.org/10.1016/s0344-0338\(97\)80104-5](https://doi.org/10.1016/s0344-0338(97)80104-5)



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.