

Loss of Blood Group Antigens in Haematolymphoid Malignancy: A Case Series from a Cancer Institute from North East India

Ranjita Sarma

Department of Transfusion Medicine, Dr B Borooah Cancer Institute, Guwahati, Assam, India.

Karabi Kuli

Department of Transfusion Medicine, Dr B Borooah Cancer Institute, Guwahati, Assam, India.

Chandana Kalita

Department of Transfusion Medicine, Dr B Borooah Cancer Institute, Guwahati, Assam, India.

Asif Iqbal

Department of Medical Oncology, Dr B Borooah Cancer Institute, Guwahati, Assam, India.

Sakshi Gupta

Department of Oncopathology, Dr B Borooah Cancer Institute, Guwahati, Assam, India.

Introduction: Altered expression of blood group antigens has been reported to be associated with both solid as well as haematological malignancies. This usually results from genetic mutation leading to incomplete as well as abnormal synthesis of antigens. Transitional loss of RBC antigens especially from ABO system is most commonly found in haematological malignancies causing blood group discrepancies and followed by reversal to its historical blood group during remission. Loss of A, B antigens has also been associated to play a role in lung, bladder and colon cancers. However not much data is available regarding haematological malignancies.

Methods and Methodology: Retrospective review of blood group discrepancy (Forward group) in patients admitted for haematolymphoid malignancy at our institute was done during a period of 3.5 years (13/3/2020 till 12/9/2023). Blood grouping done by Column agglutination test were repeated by conventional tube method. Further confirmation of the group was done by saliva test and adsorption elution test as indicated. The detail history of the patient was collected from the hospital records and analysed.

Results: A total of 7 patients presented with either loss or decrease in antigen expression. Most common diagnosis was Acute Myeloid Leukaemia (AML) with FLT3 mutation and most common antigen affected was A.

Conclusion: Haematolymphoid malignancies has been associated with loss of blood group antigens causing group discrepancy. Blood transfusion is an integral part of the supportive care in these patients. Hence, proper work up of blood group discrepancy cases should be done and recorded so as to prevent delay in blood transfusion.

Introduction

The ABO system is the most important of all blood groups in transfusion practice. A, B and H antigens are the small carbohydrate epitopes present in the glycoproteins and glycolipids of erythrocytes, endothelial cells and most epithelial cells [1]. Blood group discrepancy is said to occur when the forward blood group does not match with the results of the reverse blood group. These can be due to conditions related to patient serum (reverse grouping), or with patient's red cells (forward grouping) or with both serum and cells [2]. Both solid and haematological malignancies have been associated to alter the A, B and H antigen expression leading to discrepancy in the forward group [3, 4]. The loss of blood group antigen was first reported by Van,

loghem.et al [5] in the year 1957. The transient loss may be due to inactivation of A/B transferases in chromosome 9 or inactivation of H transferases in chromosome 19 [6, 7]. It may also have a diagnostic and prognostic implications in many cancer patients. They are more commonly seen in haematological malignancy like leukaemia, lymphoma etc [2]. Blood transfusion is an integral part of the supportive care of haematological patients and for that accurate blood group of the patient must be known. We have very limited data regarding such cases from North Eastern population of India. Hence all such rare blood group discrepancies must be solved and properly documented so as to avoid any delay in blood transfusion and to prevent haemolytic transfusion reactions.

Here we present a case series on the ABO discrepancies due to decrease or loss of expression of blood group antigens in haematolymphoid malignancy.

Materials and Methods

Blood centre of our Institute received seven samples from department of medical oncology with decrease in antigen expression in the forward blood group over a period of 3.5 years (13/3/2020 till 12/9/2023). In all seven cases, 5ml of blood sample was collected in a EDTA vial and blood grouping was done using Column agglutination test (CAT) by BIORAD. If any discrepancy was found between forward and reverse grouping, first clerical error is ruled out and then detail clinical history of the patient is taken from the hospital records. In forward blood group, when there is a loss or decrease expression of antigen by CAT, a repeat blood grouping is done by conventional tube test with monoclonal Anti-A, Anti-B and Anti-D (Tulip diagnostics) after incubation at 4°C for 30 minutes and reverse grouping is done using in-house prepared A, B and O pooled cells. If blood group is still inconclusive, saliva inhibition test is done followed by adsorption and elution studies in few cases. Each time the patient visited the cancer institute, the blood group was repeated by blood centre as all of them were on regular blood transfusion. All the procedures were done following standard technical manuals [8] as outlined in flowchart (Figure 1) and departmental standard operating procedure.

Figure 1. Flowchart Showing Test Sequence for Solving ABO Discrepancy.

Results

Observations

A total of 7 cases were reported with loss or decreased antigen expression in forward grouping (Table 1).

	Forward Group			Reverse Group			Saliva test	Adsorption-Elution study
	Anti A	Anti B	Anti D	A Cells	B Cells	O Cells		
Case 1	Neg	Neg	4+	Neg	4+	Neg	A, H	Not done
Case 2	Neg	Neg	Neg	Neg	4+	Neg	Not done	A antigen present
Case 3	Neg	Neg	4+	Neg	3+	Neg	A, H	Not done
Case 4	Neg	Neg	4+	4+	Neg	Neg	B, H	Not done
Case 5	Weak positive	Neg	Neg	Neg	4+	Neg	None	A antigen present.
Case 6	Neg	Neg	4+	Neg	3+	Neg	None	Not done
Case 7	Neg	Neg	4+	Neg	3+	Neg	A, H	Inconclusive

Table 1. Blood Group Detection by Various Methods.

Loss of expression of antigen was highest with A blood group (5 cases) followed by B group in one case whereas weak expression of antigen was seen in one case. Over all 85.5% (six) cases were Blood group A. On incubation at 4°C for 30 minutes, only single sample i.e. (Case 5) showed increase in the strength of reaction to 1+. Rest all cases still showed no positive reactions. Saliva test could be done in 6 cases out of which 4 were secretors (66.6%). Adsorption elution was done in three cases out of which two cases showing blood group as A and one was inconclusive.

On subsequent follow up, historical blood group started appearing within (1-2) months of starting induction therapy (Table 2).

	Age	Gender	Diagnosis	Molecular Study	Drug	Forward Group		Appearance of original group
						Before	After	
Case1	28yr	Male	AML	FLT3 mutation	Cytarabine	A +ve	O +ve	60 days
Case 2	8 yr	Male	AML	FLT 3 Mutation	Cytarabine	A -ve	O -ve	37 days
Case 3	36yr	Male	AML	Normal cytogenetics	Chemotherapy not started	A +ve	O +ve	30 days
Case 4	68yr	Female	AML	CEBPA gene	Decitabine	B +ve	O +ve	45 days
Case 5	32yr	Male	AML	FLT3 mutation	Cytarabine	A +ve	Weak expression of Antigen A	42 days
Case 6	56yr	Female	T-NHL Grade IV	CD20, CD3>>CD 20	CVP therapy*	A +ve	O +ve	32 days
Case 7	34yr	Malea	CML	t (9/22)	Dasatinib	A +ve	O +ve	Lost to follow up

Table 2. Patient Characteristics and Follow up.

*CVP (Cyclophosphamide +Vincristine +Prednisolone); AML, Acute Myeloid Leukaemia; CML, Chronic Myeloid Leukaemia; T-NHL, T cell - Non Hodgkin Lymphoma

During remission, the forward blood group first showed mixed field reaction by CAT followed by gradual increase in the expression of the corresponding antigens. However complete reversal to historical blood group could be seen in 3 cases and remaining was lost to follow up. Males were affected more than females and the most common diagnosis was AML in 5 cases followed by one case each of Chronic myeloid leukaemia (CML) and T-cell Non -Hodgkin Lymphoma (T-NHL). The most mutation was FLT3 seen in all AML cases. In majority of the cases (six) the discrepancy was found during the course of the treatment and in one case (Case 3) it was present before the treatment started.

Discussion

In our case series, blood group A was most commonly affected (85.7%) as seen in multiple case reports [9-11]. Study done by Abegaz SB et al [12] has shown significant decrease in expression of A, B or H antigens between 17% and 37% in leukaemia patients when compared to healthy controls. Majority of our patients presented with AML (71%) similar to other case reports [11, 13]. We also had one case each of CML and T-NHL. Loss of blood group antigens in CML was also

reported by Chenna D et al [11] but no literature could be found related to T-NHL. In one study by Xiros N et al, there was a change in the blood group of a myelodysplastic syndrome patient following blastic transformation to AML [10]. Among the AML patients, most common mutation seen was FLT3 and during induction phase most common drug used was cytarabine. Prakash S et al also reported a similar case of AML on induction therapy with cytarabine but the mutation was in CEBPA genes [9]. Saliva test done to confirm the blood group showed 66.6% as secretors which was very less compared to other Indian studies [14, 15]. It might be due to small sample size of the study or due to disease itself. Historical blood group antigens started appearing between 30 to 60 days after first cycle of induction chemotherapy and complete reversal of blood group could be seen in three cases. During this transition period, initially we observed mixed field reactions in CAT followed by complete reversal of blood group to the original one similar to a case reported by Prakash S et al [9]. In our study, majority of the patients were on chemotherapy (85.7%) and in one patient the chemotherapy was not started. This may be due to loss of antigen expression during the course of disease progression or during chemotherapy and it reappeared during remission period. Multiple causes like inhibitory factor related to antigen-antibody binding or deficiency in transferase synthesis activity or an abnormal distribution or density of antigen sites in the RBC membrane have been linked to decrease in antigen expression [16]. There is paucity of data and research materials related to blood group discrepancy in cancer patients. We could not establish the exact cause of the discrepancy at the molecular level. In this case series, we could however highlight that the loss of blood group antigens was most common in AML patients and blood group A being the most commonly affected group. Hence, we need further molecular and genetic studies to establish the basis for such hypothesis and its implication in disease progression and prognosis.

In conclusion, Haematolymphoid malignancy may show altered expression of blood group antigens. All cases of blood group discrepancy should be properly evaluated and documented. This not only prevents the delay in providing safe blood in case of emergency but also gives us a gist regarding underlying disease condition. Further research and development programme has to be initiated to know the association between loss of antigen expression and cancer biology.

Acknowledgements

The authors acknowledge Mr Hemanta Deka and Mr Arup Baishya, Blood Centre Technician, Dr B Borooah Cancer Institute, Guwahati, Assam, India.

References

References

1. Laura. Carbohydrate blood groups. In: Rossi's principles of transfusion medicine. 5th ed. West Sussex: John Wiley and Sons Limited. 2016;159-175.
2. Denise M. Harmening. ABO blood group systems. In: Modern Blood Banking and Transfusion Practices. 6th. Philadelphia: F.A. Davis Company. 2012;119-148.
3. Klein, Harvey G, David J. Anstee. Immunology of red cells. In: Mollison's blood transfusion in clinical medicine. 12th edition. 2014;53-117.
4. Bianco T, Farmer BJ, Sage RE, Dobrovic A. Loss of red cell A, B, and H antigens is frequent in myeloid malignancies. *Blood*. 2001; 97(11)[DOI](#)
5. Van Loghem JJ, Dorfmeier H, Van Der Hart M. Two A antigens with abnormal serologic properties. *Vox Sanguinis*. 1957; 2(1)[DOI](#)
6. Crookston MC. Anomalous ABO, H, Ii phenotypes in disease. In: Garratty G, editor. Blood Group Antigens and Disease. Arlington, VA: American Association of Blood Banks. 1983;67-84.



7. Hoogstraten B, Rosenfield RE, Wasserman LR. Change of ABO blood type in a patient with leukemia. *Transfusion*. 1961; 1^{DOI}
8. Technical manual, 20th edition: methods and appendices. Method 2.7,2.8. AABB website. <https://www.aabb.org/aabb-store/resources/technical-manual-methods>.
9. Prakash S, Mohapatra S, Bhagavathi MS, Das N, Krushna Ray G, Mukherjee S. Loss and Reappearance of A Antigen After Chemotherapy Leading to Blood Group Discrepancy in Acute Myeloid Leukemia: A Case Report. *Laboratory Medicine*. 2021; 52(5)^{DOI}
10. Xiros N, Northoff H, Anger B, Heit W, Heimpel H. Blood group change in a patient with blastic transformation of a myelodysplastic syndrome. *Blut*. 1987; 54(5)^{DOI}
11. Chenna D, Mohan G, Reddy VR, Shastry S. The disappearance of blood group antigens: A clue to the clinical diagnosis of leukemia. *Transfusion and Apheresis Science: Official Journal of the World Apheresis Association: Official Journal of the European Society for Haemapheresis*. 2019; 58(1)^{DOI}
12. Abegaz SB. Human ABO Blood Groups and Their Associations with Different Diseases. *BioMed Research International*. 2021; 2021^{DOI}
13. Nambiar RK, Narayanan G, Prakash NP, Vijayalakshmi K. Blood group change in acute myeloid leukemia. *Proceedings (Baylor University Medical Center)*. 2017; 30(1)^{DOI}
14. Rajawat G, Ramalingam K, Pareek R, Singh G, Narula H, Aggarwal A. Assessment of Salivary ABO Blood Group Antigens and Secretor Status in Sriganganagar, Rajasthan: A Correlational Analysis of 300 Samples. *Cureus*. 2023; 15(4)^{DOI}
15. Metgud R, Khajuria N, Mamta N, Ramesh G. Evaluation of the Secretor Status of ABO Blood Group Antigens in Saliva among Southern Rajasthan Population Using Absorption Inhibition Method. *Journal of clinical and diagnostic research: JCDR*. 2016; 10(2)^{DOI}
16. Ewald DR, Sumner SCJ. Blood type biochemistry and human disease. *Wiley Interdisciplinary Reviews. Systems Biology and Medicine*. 2016; 8(6)^{DOI}