

# Efficacy of Extracorporeal Irradiation in Primary Malignant Bone Tumours: A Tertiary Cancer Centre Experience

*Gunaseelan k*

*Dilip Kumar Patro*

Professor Department of Orthopaedics JIPMER, India.  
3Senior Resident, Department of Orthopaedics, JIPMER, India.

*Arun Lal*

Senior Resident, Department of Orthopaedics, JIPMER, India.

*Kannan P*

Senior Resident, Department of Radiation Oncology, JIPMER, India.

*Biswajit D*

Additional Professor, Department of Medical Oncology, JIPMER, India.

*Vijayaprabhu N*

Assistant Professor, Medical Physics, Department of Radiotherapy, JIPMER, India.

**Background:** Limb-sparing approaches involving a combination of surgery, chemotherapy, and radiotherapy are preferred in primary malignant bone tumours whenever possible. The limb function can be maintained by performing reconstruction either using a prosthesis or bone graft. Extracorporeal irradiation is a novel and exciting technique in which the restoration is done using the patient's own tumour bone. In this article, we present the local control, complications, and functional outcome of 49 cases of primary malignant bone tumours treated using the technique of extracorporeal irradiation.

**Materials and methods:** 49 patients with primary malignant bone tumours were treated with extracorporeal irradiation between the year 2007 to 2015. En-bloc resection of the tumour and the involved bone along with soft tissues was performed about four weeks after the completion of three to four cycles of neoadjuvant chemotherapy. The bone segment was irradiated with 6MV linear accelerator to a dose of 50Gy. After extracorporeal irradiation, the bone was re-implanted using fixation devices. Local control, relapse-free survival, complications and functional outcome were studied.

**Results:** Out of 49 patients, there were 33 males and 16 females. The median age was 19 years. Femur and tibia were the most common tumour locations with osteosarcoma being the most common histology (74%). The local control rate was 94%. The 8 year local recurrence-free survival and distant metastasis-free survival were 89% and 84% respectively. The overall complication rate was 20%. Infection was the most frequent complication, observed in 46% of the patients who developed complications. Out of 49 patients, 40 had reported a good range of movements.

**Conclusion:** Extracorporeal irradiation and re-implantation is a convenient alternative to prosthesis from a health economics point of view, provided severe complications are managed and avoided so that functional results can be achieved.

---

## Introduction

Primary malignant bone tumours are common in children and adolescents because of active skeletal growth. About sixty percentage cases occur between 10 and 20 years of age [1]. The standard management approach for malignant bone tumours is multidisciplinary, especially when

there is a concern for limb salvage. The treatment of malignant bone tumours has undergone a drastic change in the last 20 years. Limb-sparing strategies are preferred whenever possible, which may involve a combination of surgery, chemotherapy, and radiotherapy [2][3][4][5][6]. The limb function can be maintained by performing reconstruction either using a prosthesis or bone graft. However, in developing countries, many patients would not be able to afford for prosthesis due to poor socioeconomic status or there may be a delay in procurement of prosthesis leading on to delay in the management. Moreover, prosthesis may not provide precise anatomic fit and can be associated with problems like loosening and instability. Recently, there has been an excessive interest in using the patient's tumour bone for reconstruction after sterilisation, to preserve the limb function. The various methods of sterilisation include autoclaving, pasteurisation, use of liquid nitrogen and extracorporeal irradiation (ECI). Extracorporeal irradiation is a novel and exciting technique. In ECI, the tumour bearing segment of bone is excised en bloc; the tumour and the soft tissues are then removed from the bone, and the bone is irradiated and reimplanted back into the body [7][8][9][10][11]. In this article, we analyse and present the results of malignant bone tumours treated using the technique of extracorporeal irradiation.

## Materials and Methods

Forty-nine histologically proved non-metastatic primary malignant bone tumours suitable for limb preservation therapy, from the year 2007 to 2015, were included in this study after tumour board discussion. The eligible patients underwent a clinical examination by a multidisciplinary team and magnetic resonance imaging of the affected limb to assess the local extent of the disease. Contrast-enhanced CT of the thorax and bone scan were done to exclude lung and bone metastases respectively. Other routine investigations included complete haemogram, renal function test, and liver function test. The patients were usually treated with neoadjuvant chemotherapy followed by surgery, ECI and reimplantation followed by adjuvant chemotherapy. Patients with osteosarcoma were given IAP [Ifosfamide 1.5g/m<sup>2</sup> day 1,2&3, Adriamycin 60mg/m<sup>2</sup> day 1, and cisplatin 100mg/m<sup>2</sup> divided over days 1-3) regimen every three weeks for six cycles. Ewings sarcoma patients were treated with alternating VAC (vincristine 1.5mg/m<sup>2</sup> day 1, Adriamycin 60mg/m<sup>2</sup> day 1, and cyclophosphamide 1200mg/m<sup>2</sup> day 1) and IE (ifosfamide 1.8g/m<sup>2</sup>/day for 5 days, etoposide 100mg/m<sup>2</sup>/day for 5 days) regimen every three weeks for a total of 16 cycles.

Neoadjuvant chemotherapy was given for three to four cycles. Surgery was performed about four weeks after the completion of neoadjuvant chemotherapy. The surgical procedure and ECI technique is summarised inb figure. During operation, the tumour bearing segment of the bone is removed en bloc along with soft tissues. The tumour and the soft tissues are then removed from the bone, which is wrapped in a sterile towel and further twice wrapped in sterile plastic bags and sent for radiation in an acrylic phantom. Sterile saline water (tissue equivalent) was poured in the phantom. The bone segment was irradiated in the radiotherapy department with 6 MV linear accelerator to a dose of 50 Gy using parallel opposed lateral beams. During the ECI, the operative site was prepared for reimplantation, and biopsy was performed at all osteotomy sites to assess the status of resection margins. After ECI, the bone was then reimplanted using fixation devices under C-arm guidance. During the postoperative period, immobilization was continued until the radiographic imaging showed evidence of union. Full weight bearing was allowed according to the clinical and radiological progress.

### Figure 1. Disease status.

The local control, relapse-free survival, complications and functional outcome were studied. The functional outcome was reported as good, fair, and poor based on the criteria used by Uyttendaele et al [12], as shown in table 1.

Criteria used to classify the functional results
--

Good
Fair
Poor

**Table 1. Assessment of orthopaedic outcome.**

## Results

49 patients with primary malignant bone tumours were treated with extracorporeal irradiation technique at our institute between 2007 and 2015. The clinical and tumour characteristics were shown in table 1. The predominant histology was osteosarcoma(74%). Other histologies included ewings sarcoma and chondrosarcoma. Majority of patients were males(67%). The age group of the patients ranged from nine to 40 years and the median age was 19 years. The long bones of the lower limbs were most frequently affected and nearly 90% of the tumors were located in femur and tibia. The median duration of follow up was 24 months (range 4-108 months).

Characteristic	Number	Percentage
<b>Gender</b>		
Male	33	67
Female	16	33
<b>Age</b>		
Median	19 years	
Range	9-40 years	
<b>Tumour location</b>		
Femur	24	49
Tibia	20	41
Humerus	2	4
Radius	2	4
Fibula	1	2
<b>Histology</b>		
Osteosarcoma	36	74
Ewings sarcoma	4	8
Chondrosarcoma	3	6
Others	6	12
<b>Follow up</b>		
Median	24 months	

Range	4-108 months
-------	--------------

**Table 2. Patient and tumour characteristics.**

Ten patients had developed recurrence either in the form of only local (two patients), distant metastases (seven patients) or both (one patient). The local control rate was 94%. The 8 year local recurrence free survival was 84%. The 8 year distant metastasis free survival was 84%. The type of complication that had occurred in the perioperative period is shown in figure.

**Figure 2. Complications.**

The overall complication rate was 20%. The most frequent complication was infection, seen in 46% of the patients who developed complications.

**Figure 3. Extracorporeal irradiation technique.**

With respect to orthopedic outcome, 40 patients had reported good range of movements as shown in figure 4. At the end of last follow up, out of 49 treated patients, 37 patients were alive, 9 patients had expired and 3 were lost to follow up.

**Figure 4. Orthopaedic outcome.**

## Discussion

Limb sparing procedures are the preferred options for malignant bone tumours whenever feasible [4]. Custom made prostheses or osteoarticular allografts can be used for reconstruction but prostheses are expensive and cannot be afforded by all patients in developing countries. Moreover, high complication rates, with 30% infection and 23% nonunion, have been reported using allografts, especially in children. Patients with graft complications had a significantly lower probability of graft survival, resulting in a worse functional outcome [7][8][13]. These problems can be overcome by the technique of using the patients own bone as autograft for reconstruction after extracorporeal irradiation. This technique of extracorporeal irradiation and reimplantation was first developed by Spira and Lubin [14]. There are only limited reports of ECI available in the literature [9][14][15][16]. ECI has several advantages over other methods of limb reconstruction. These include precise anatomic fit, preservation of joint mobility, avoidance of limb length discrepancy and problems associated with allograft, such as dependence on a bone bank, graft rejection, and the risk of viral transmission such as the human immunodeficiency and hepatitis C viruses.

Extracorporeal irradiation as an useful, cost effective alternative during limb salvage in bony malignancies had been reported by various studies. The studies by Davidson et al [15], and Poffyn et al [16]. were the largest in the literature of ECI that had treated 50 or more patients. Ours is the next largest study that had included 49 patients of primary malignant bone tumours managed with ECI. Davidson et al [15]. in study treated 50 patients with extracorporeal irradiation and reimplantation and found four recurrences at a mean follow up of 38 months. Poffyn et al [16]. retrospectively reviewed 107 patients with 108 malignant or locally aggressive bone tumours treated between 1978 and 2009 by extracorporeal irradiation with 300 Gy to eradicate the tumour,

and reimplantation of the bone as an orthotopic autograft. No local recurrences were detected in the irradiated grafts. In the study of 12 patients with Ewings sarcoma by Puri et al [17], six patients were free of disease and six patients had died from the disease at the time of last review and there were two recurrences around the operative site but away from the irradiated graft. Sharma et al [18]. treated 12 patients of primary malignant bone tumours with 50 Gy ECI dose and found 3 local recurrences at a median follow up of 22 months. The 2 year local recurrence free survival was 73%. In our study, ten patients had developed recurrence either in the form of only local (two patients), distant metastases( seven patients) or both (one patient) at a median follow up of 24 months. The local control rate was 94%. The 8 year local recurrence free survival was 89%. At the end of last review, out of 49 treated patients, 37 patients were alive, 9 patients had expired and 3 were lost to follow up. Thus, the local control rate and recurrences were varying among the studies, which could be attributed to the small sample size and varying doses of radiation used for ECI.

The optimal dose of ECI has not been determined. The radiation doses varying from 50 to 300 Gy had been used in the literature. Chen et al [19], Uyttendaele et al [12], and Poffyn et al [16]. used doses of 300 Gy. In our study, we delivered midplane dose of 50Gy in one fraction using Source to axis distance(SAD) technique with parallel opposed lateral beams. Hong et al [9], and Davidson et al [15], delivered doses of 50 Gy similar to our study. The risk of wound infection and other complication in various ECI series had been reported to be upto 8-17% [17][20]. Sharma et al [18], found wound infection rate of 14% ( 2 out of 14 patients). In our study, the infection rate was 12% which is similar to that reported in the literature. The literature review of various local complications is shown in table 3.

	Hong et al.(9)	Sys et al.(21)	Uyttendaele et al.(12)	Davidson et al.(15)	JIPMER
	N=16	N= 15	N=15	N=50	N=49
Infection	1	3	2	3	6
Nonunion	0	1	1	9	1
Fracture	1	0	1	3	4
Amputation	1	0	3	0	0
Reoperation	0	1	5	17	1

**Table 3. Literature review of local complications.**

The limitation of this study is that it is not a randomized study to establish extracorporeal irradiation and reimplantation as the standard technique in bone tumours when limb salvage is concerned. Even in the literature, only retrospective and non randomised studies are available. Hence, in the future, it would be worth doing a randomised study comparing extracorporeal irradiation to prostheses.

In conclusion, limb salvage approach involving multidisciplinary team is the current recommended treatment of choice for primary malignant bone tumours whenever feasible. Extracorporeal irradiation is a convenient alternative to prosthesis since affordability is a matter of concern in developing countries. If serious complications can be managed or avoided, functional results can be achieved. Thus, from a health economics point of view, ECI is a relatively low-cost technique compared with the high cost of prostheses. However, the most important determinants of the feasibility of ECI include tumour location and extent, and appropriate selection of patients.

## References

## References

1. Eyre Rachel, Feltbower Richard G., Mubwandarikwa Emmanuel, Eden Tim O.B., McNally

- Richard J.Q.. Epidemiology of bone tumours in children and young adults. *Pediatric Blood & Cancer*. 2009; 53(6)[DOI](#)
2. Bernstein M.. Ewing's Sarcoma Family of Tumors: Current Management. *The Oncologist*. 2006; 11(5)[DOI](#)
3. Donaldson SS, Torrey M, Link MP, Glicksman A, Gilula L, Laurie F, et al. A multidisciplinary study investigating radiotherapy in Ewing's sarcoma: end results of POG #8346. *Pediatric Oncology Group. Int J Radiat Oncol Biol Phys*. 1998 ; 42(1):125-135.
4. Gebhardt MC, Eyre R, Feltbower RG, Mubwandarikwa E, Eden TO, McNally RJ. What's new in musculoskeletal oncology. *Int J Radiat Oncol Biol Phys*. 2002; 84-A(4):694-701.
5. Hogendoorn P. C. W., Athanasou N., Bielack S., De Alava E., Tos A. P. D., Ferrari S., Gelderblom H., Grimer R., Hall K. S., Hassan B., Hogendoorn P. C. W., Jurgens H., Paulussen M., Rozeman L., Taminiau A. H. M., Whelan J., Vanel D.. Bone sarcomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology*. 2010; 21(Supplement 5)[DOI](#)
6. Jaffe Norman, Patel Shreyaskumar R., Benjamin Robert S.. Chemotherapy in Osteosarcoma: Basis for Application and Antagonism to Implementation; Early Controversies Surrounding its Implementation. *Hematology/Oncology Clinics of North America*. 1995; 9(4)[DOI](#)
7. Cannon SR. Massive prostheses for malignant bone tumours of the limbs. *J Bone Joint Surg Br*. 1997; 79(3):497-506.
8. Hejna Michael J., Gitelis Steven. Allograft prosthetic composite replacement for bone tumors. *Seminars in Surgical Oncology*. 1997; 13(1)[DOI](#)
9. Hong Angela, Stevens Graham, Stalley Paul, Pendlebury Susan, Ahern Verity, Ralston Anna, Estoesta Edgar, Barrett Ian. Extracorporeal irradiation for malignant bone tumors. *International Journal of Radiation Oncology\*Biophysics*. 2001; 50(2)[DOI](#)
10. Khattak Mujahid Jamil, Umer Masood, Haroon-ur-Rasheed, Umar Mohammad. Autoclaved Tumor Bone for Reconstruction. *Clinical Orthopaedics and Related Research*. 2006; 447[DOI](#)
11. Singh Vivek, Nagalingam Janarthan, Saad Marniza, Pailoor Jayalakshmi. Which is the best method of sterilization of tumour bone for reimplantation? a biomechanical and histopathological study. *BioMedical Engineering OnLine*. 2010; 9(1)[DOI](#)
12. Uyttendaele D, De Schryver A, Claessens H, Roels H, Berkvens P, Mondelaers W. Limb conservation in primary bone tumours by resection, extracorporeal irradiation and re-implantation. *The Journal of Bone and Joint Surgery. British volume*. 1988; 70-B(3)[DOI](#)
13. Mankin Henry J., Gebhardt Mark C., Jennings L. Candace, Springfield Dempsey S., Tomford William W.. Long-Term Results of Allograft Replacement in the Management of Bone Tumors. *Clinical Orthopaedics and Related Research*. 1996; 324[DOI](#)
14. Spira E, Lubin E. Extracorporeal irradiation of bone tumors. A preliminary report. *Isr J Med Sci*. 1968 Sep; 4(5):1015-1019.
15. Davidson A. W., Hong A., McCarthy S. W., Stalley P. D.. En-bloc resection, extracorporeal irradiation, and re-implantation in limb salvage for bony malignancies. *The Journal of Bone and Joint Surgery. British volume*. 2005; 87-B(6)[DOI](#)
16. Poffyn Bart, Sys Gwen, Mulliez Alexander, Van Maele Georges, Van Hoorebeke Luc, Forsyth Ramses, Uyttendaele Dirk. Extracorporeally irradiated autografts for the treatment of bone tumours: tips and tricks. *International Orthopaedics*. 2010; 35(6)[DOI](#)
17. Puri Ajay, Gulia Ashish, Agarwal MG, Jambhekar NA, Laskar S. Extracorporeal irradiated tumor bone: A reconstruction option in diaphyseal Ewing's sarcomas. *Indian Journal of Orthopaedics*. 2010; 44(4)[DOI](#)
18. Sharma DN, Rastogi S, Bakhshi S, Rath GK, Julka PK, Laviraj MA, Khan SA, Kumar A. Role of extracorporeal irradiation in malignant bone tumors. *Indian Journal of Cancer*. 2013; 50(4)[DOI](#)
19. Chen W. M., Chen T. H., Huang C. K., Chiang C. C., Lo W. H.. Treatment of malignant bone tumours by extracorporeally irradiated autograft-prosthetic composite arthroplasty. *The Journal of Bone and Joint Surgery*. 2002; 84(8)[DOI](#)
20. Araki Nobuhito, Myoui Akira, Kuratsu Sigeyuki, Hashimoto Nobuyuki, Inoue Takehiro, Kudawara Ikuo, Ueda Takafumi, Yoshikawa Hideki, Masaki Norie, Uchida Atsumasa. Intraoperative Extracorporeal Autogenous Irradiated Bone Grafts in Tumor



Surgery. *Clinical Orthopaedics and Related Research*. 1999; 368 [DOI](#)