

Factors Favoring Long Term Survival in Patients with Stage IV Epithelial Ovarian Cancer: An Institutional Research

Hafiz Abubakar Sarwar

Jhanzeb Iftikhar

Department of Medical Oncology, Shaukat Khanum memorial cancer hospital and Research center, Lahore, Pakistan.

Samia Yasmeen

Department of Medical Oncology, Shaukat Khanum memorial cancer hospital and Research center, Lahore, Pakistan.

Fareeha Sheikh

Department of Medical Oncology, Shaukat Khanum memorial cancer hospital and Research center, Lahore, Pakistan.

Fatima Ali

Department of Medical Oncology, Shaukat Khanum memorial cancer hospital and Research center, Lahore, Pakistan.

Aamir Ali Syed

2) Department of Surgical Oncology, Shaukat Khanum memorial cancer hospital and Research center, Lahore, Pakistan.

Neelam Siddiqui

Department of Medical Oncology, Shaukat Khanum memorial cancer hospital and Research center, Lahore, Pakistan.

Objective: Our aim was to identify factors favoring long term survival in patients presenting with stage IV epithelial ovarian cancer.

Methods: We did retrospective analysis of thirty patients with stage IV epithelial ovarian cancer diagnosed and treated at Shaukat Khanum Memorial Cancer Hospital, Lahore, Pakistan from 2006 to 2013. Patient's demographics, clinical data and histopathology were abstracted from cancer registry department of our hospital. Chi-square test was used to find the association between clinic-pathological variables and long term survival.

Result: All patients received chemotherapy and surgery as per ovarian cancer guidelines. Of the thirty patients, eleven patients survived greater than four years median survival was recorded as thirty five months. Absence of co-morbidities and good performance status indicated good results of therapy however did not have statistically significant impact on survival. Higher CA-125 at presentation i.e. >1000 (normal range : <21 U/ml), response to initial chemotherapy, interval cytoreductive surgery and complete response after induction therapy were significantly associated with long term survival ($P < 0.05$).

Conclusion: Prognosis of patients presenting with stage IV epithelial ovarian cancer remains poor. Very high values of CA-125 (>1000) at presentation, response to initial chemotherapy, interval surgical resection and complete remission after induction therapy, appear to be significant prognostic factors for long term survival. Further studies exploring molecular profiling and immunological factors are warranted.

Introduction

Ovarian cancer accounts for 3% of all cancers among women and is the 5th leading cause of cancer related death world wide [1]. Epithelial ovarian cancer (EOC) accounts for 90% of all malignant tumors of ovary and these are adenocarcinomas. EOC are further classified into Type I and Type II

tumors [2][3]. Type I tumors are considered low grade and include endometrioid, mucinous and clear cell histologies and they carry mutations of BRAF, KRAS and PTEN with microsatellite instability. While type II tumors are high grade serous and carcinosarcomas which harbor mutations in BRCA1, BRCA2 and p53 [4].

Ovarian cancer usually presents in postmenopausal women with history of abdominal pain and distension. Seventeen percent of the patients present with stage IV disease, where 5-years survival rate remains less than 10% [5][6]. Currently with advanced treatment modalities, a significant number of patients achieve complete biochemical and radiological response. However patients with stage IV disease tend to relapse within eighteen months after treatment completion [3]. Among the relapsed patients, in whom tumor remains sensitive to platinum based chemotherapy, outcomes are good as compared to those whose disease becomes platinum refractory in subsequent relapses [3].

Approximately 295,000 women developed ovarian cancer every year world wide out of which 90% are epithelial ovarian cancers (1). In Pakistan, its incidence is also rising by 1-2% every year and various studies conducted in Pakistan report it as among the ten most common cancers in Pakistan [7][8]. Cress RD et al., evaluated characteristics of long term survival in 3,582 patients of ovarian cancer at all stages of presentation and found that younger age, early stage, low grade and non-serous histology are predictors of long term survival, but some women with high risk cancer lived longer [9]. Hamilton CA et al., conducted a study on 3010 patients with advanced stage epithelial ovarian cancer and found that 195 patients survived greater than 10 years. Out of these patients, the number of stage IV patient were very low and the outcomes were worse. They revealed that lower CA-125 value, absence of ascites, stage at presentation and R0 resection had significant influence on long term survival [10]. It is observed that there is no significant survival data available for Pakistani population due to lack of studies. Secondly majority of existing literature shows combined data of stage III and stage IV so we specifically studied stage IV patients in this study.

Materials and Methods

The present time based research was conducted in SKMCH&RC. All patients who presented with stage IV epithelial ovarian cancer from 2006 to 2013 were included. Thirty patients with newly diagnosed stage IV EOC were analyzed. Patient's baseline characteristics including age, comorbidities, ECOG-performance status and BMI were recorded. Details of therapy including surgical procedure underwent obtained. All of the patients received neoadjuvant platinum containing regimens as per ovarian cancer protocol. CA-125 was recorded at baseline, after completion of treatment and on subsequent follow-ups. The histological type, site of metastasis and ascites were also recorded.

Statistical analysis

SPSS software (version 23.0; SPSS, Chicago, IL, USA) was used for statistical analysis of the data. Mean \pm standard deviation was used for continuous variables while frequencies and percentages used for categorical variables. Chi square test or fisher's exact test (when necessary) were used to compare categorical variable. The Kaplan-Meier method was used to estimate survival as a function of time, and survival differences were analyzed by the log-rank test. Statistical significance was defined as a two-tailed p-value 0.05. Approval for study was taken from our Institutional Review Board (IRB).

Results

Table 1 presents the demographic and clinical characteristics of 30 stage IV ovarian cancer patients with a mean age and standard deviation of 50 ± 13 years. All the patients were females with mean body mass index and standard deviation of 27 ± 6.0 . In addition, more than a half (57%)

of the respondents had no comorbidity and (23%) had multiple comorbidities. Additionally, majority (63%) of the patients were died on last follow-up and the median survival was survival was found thirty five months. Among patients who survived, half of them had ECOG performance status of 2 (meaning that patients were ambulatory and capable of all self-care but unable to carry out any work activities; up and about more than 50% of waking hours) Stratification of age, body mass index, comorbidity and ECOG performance status with respect to status (alive and death) is shown in Table 1.

Variables	Categories	Total = N (%)	Alive 11 (36.7%)	Death 19 (63.3%)	p-value
Age (years)	Mean \pm SD*	49.97 \pm 12.95	48.55 \pm 6.96	50.79 \pm 15.53	0.66
Body Mass Index	Mean \pm SD*	27.33 \pm 5.70	26.89 \pm 5.34	27.58 \pm 6.03	0.75
Comorbids	No	17 (56.7%)	6 (35.3%)	11 (64.7%)	0.71
	Single	6 (20.0%)	3 (50.0%)	3 (50.0%)	
	Multiple	7 (23.3%)	2 (28.6%)	5 (71.4%)	
ECOG-PS	1**	12 (40.0%)	5 (41.7%)	7 (58.3%)	0.38
	2**	15 (50.0%)	6 (40.0%)	9 (60.0%)	
	3**	3 (10.0%)	0 (0.0%)	3 (100.0%)	

Table 1: Descriptive statistics (Demographics and Clinical characteristics).

***standard deviation, 1** = Ambulatory, 2** = In bed <50%, 3** = In bed >50%**

Furthermore, Table 2 presents the pathological characteristics of patients with an overall mean CA-125 value and standard deviation of 4100 \pm 954. Majority of the patients (63%) had serous histopathology and more than half (53%) of the patients had M-1 site of metastasis. Moreover, surgical procedure was performed in (70%) of the patients and more than half of patients showed complete response to chemotherapy as shown in Table 2. Therefore, overall survival was found statistically significant (p-value 0.002 using chi-square test) among patients who showed complete response to initial therapy as shown in Table 2.

Variables	Categories	N (%)	Alive 11 (36.7%)	Death 19 (63.3%)	p-value
CA-125	Mean \pm SE*	4100.17 \pm 954.38	5165.10 \pm 1557.34	3425.32 \pm 1211.62	0.36
Histopathology	Adeno unspecified	3 (10.0%)	0 (0.0%)	3 (100.0%)	0.64
	Serous	19 (63.3%)	8 (42.1%)	11 (57.9%)	
	Mucinous	2 (6.7%)	1 (50.0%)	1 (50.0%)	
	Others	6 (20.0%)	2 (33.3%)	4 (66.7%)	
Site of Metastasis	M-1	16 (53.3%)	5 (31.2%)	11 (68.8%)	0.71
	M-2	14 (46.7%)	6 (42.9%)	8 (57.1%)	
Ascites	Yes	26 (84.7%)	9 (34.6%)	17 (65.4%)	0.61
	No	4 (13.3%)	2 (50.0%)	2 (50.0%)	
Number of Chemotherapy cycles	Mean \pm SD**	22.0 \pm 21.0	34.91 \pm 28.61	14.47 \pm 10.20	0.01

Surgical Resection	Surgery Performed	21 (70.0%)	8 (38.1%)	13 (61.9%)	1.00
	No surgery performed	9 (30.0%)	3 (33.3%)	6 (66.7%)	
Response	Complete Response	16 (53.3%)	10 (62.5%)	6 (37.5%)	0.002
	No Complete Response	14 (46.7%)	1 (7.1%)	13 (92.9%)	
Survival Time (days)	Mean \pm SD	1197 \pm 789.94	2091.10 \pm 111.36	680.80 \pm 91.69	0.001

Table 2: Descriptive statistics (Pathological characteristics).

*standard error; ** standard deviation

Therefore, Overall survival was compared between complete response and no complete response. There was statistically significant difference ($p=0.001$ using independent sample t-test) between the two groups as shown in Figure 1.

Figure 1. Overall Survival was Compared between Complete Response and no Complete Response.

Discussion

In our study, 70% stage IV ovarian cancer patients received neo-adjuvant platinum based chemotherapy followed by surgery as per ovarian cancer guidelines. Among thirty patient twenty one (70%) patients underwent cytoreductive surgery while nine patients (30%) were not fit enough to undergo surgery. The outcomes were comparable for both groups in terms of long term survival. (10) Hamilton CA et al, similar results describing that cytoreductive surgery did not influence the outcomes and 15% of patients survived more than 10 years. Surprising results were found in a study performed by Cress RD et al., that one third of patients with stage III and IV disease and serous histology showed long term survival of more than 10 years. The possible reason for this finding was improved surgical techniques with no or minimal residual disease and concomitant use of intraperitoneal and adjuvant chemotherapy and the possibility of somatic mutations in patients with same histology (9).

A contrasting result was found regarding CA-125 level values. In our study the impact of CA-125 levels acts differently. High values of CA-125 at presentation were linked to long term survival as compared to previous studies, though it was not statistically significant ($p=0.36$). The reason for this finding might be ethnic variation in our region. Existing literature showed that lower pretreatment CA-125 values is associated with long term survival (11).

Tumor histology is also an important predictor of survival. (12-14) In our study population, the most common histological type was serous (63%) and remaining histologies were mucinous (7%), adeno-unspecified (10%) and others were (20%). None of the histological subtype was associated with statistically significant long term survival ($p\text{-value}=0.64$).

In sixteen patients (53%) the primary site of metastasis was pleural effusion (M1) while in fourteen patients (47%) metastatic sites were other than pleura (M2). Among M2 metastasis six patients (43%) had better long term survival of more than four years. In this study, twenty six (85%)

patients had ascites at the time of presentation while four patients (13%) had no ascites.

It was found that the eleven patients (37%) who responded and tolerated neoadjuvant platinum based chemotherapy showed long term survival of more than four years and the results were also statistically significant (p-value= 0.01) as compared to 19 patients (63%) whose disease was not responsive to chemotherapy and could received (14 ± 10 number of chemotherapy cycles) less number of chemotherapy cycles. The results were also significant. Melamed A, et al., and Rosen B, et al., showed that neoadjuvant chemotherapy for advance ovarian cancer has survival benefits. (15, 16)

One of the most important predictor of long term survival found in this study was achievement of complete radiological and biochemical response after neoadjuvant chemotherapy, surgery and adjuvant chemotherapy(p-value= 0.02). In this study group 16 (53%) patients achieved complete response. Among them 10(63%) patients showed long term survival of more than four years while 6 (38%) patient showed long term survival of less than four years. While 14 (47%) patients could not achieve a complete response. Among them only one patient (7%) lived for more than 4 years while 13(93%) patients survived less than four years. P value (0.002) also showed that there is an association between response status and long term survival.

In conclusion, advanced stage ovarian cancer is still a fatal cancer with poor outcomes. Achieving complete response after neoadjuvant chemotherapy, cytoreductive surgery adjuvant chemotherapy has significant prognostic impact on patient's long term survival. Further studies looking for patterns of gene expressions associated with long term survival, molecular profiling and immunological factors are needed.

Acknowledgements

I am highly indebted to my respected mentor Dr. Neelam Siddiqui, Medical Oncologist at Shaukat Khanum Memorial Cancer Hospital and Research center, Lahore, Pakistan for her valuable guidance, encouragement and constant supervision during the accomplishment of this task.

References

References

1. International Agency for Research on Cancer. *Global cancer statistics.2018*.http://globocan.iarc.fr/Page/fact_sheets_population.aspx(accessed June 6, 2019).
2. Shih Ie-Ming, Kurman Robert J.. Ovarian Tumorigenesis. *The American Journal of Pathology*. 2004; 164(5)[DOI](#)
3. Jayson Gordon C, Kohn Elise C, Kitchener Henry C, Ledermann Jonathan A. Ovarian cancer. *The Lancet*. 2014; 384(9951)[DOI](#)
4. Banerjee S., Kaye S. B.. New Strategies in the Treatment of Ovarian Cancer: Current Clinical Perspectives and Future Potential. *Clinical Cancer Research*. 2013; 19(5)[DOI](#)
5. McGuire William P., Hoskins William J., Brady Mark F., Kucera Paul R., Partridge Edward E., Look Katherine Y., Clarke-Pearson Daniel L., Davidson Martin. Cyclophosphamide and Cisplatin Compared with Paclitaxel and Cisplatin in Patients with Stage III and Stage IV Ovarian Cancer. *New England Journal of Medicine*. 1996; 334(1)[DOI](#)
6. Ozols Robert F., Bundy Brian N., Greer Benjamin E., Fowler Jeffrey M., Clarke-Pearson Daniel, Burger Robert A., Mannel Robert S., DeGeest Koen, Hartenbach Ellen M., Baergen Rebecca. Phase III Trial of Carboplatin and Paclitaxel Compared With Cisplatin and Paclitaxel in Patients With Optimally Resected Stage III Ovarian Cancer: A Gynecologic Oncology Group Study. *Journal of Clinical Oncology*. 2003; 21(17)[DOI](#)

7. Saeed S, Akram M. EPITHELIAL OVARIAN CANCER. *The Professional Medical Journal*. 2012; 19(01):040-045.
8. Bhurgri Yasmin, Bhurgri Asif, Hassan Sheema H., Zaidi S.H.M., Rahim Abdul, Sankaranarayanan Rengaswamy, Parkin D. Maxwell. Cancer incidence in Karachi, Pakistan: First results from Karachi Cancer Registry. *International Journal of Cancer*. 2000; 85(3)[DOI](#)
9. Cress Rosemary D., Chen Yingjia S., Morris Cyllene R., Petersen Megan, Leiserowitz Gary S.. Characteristics of Long-Term Survivors of Epithelial Ovarian Cancer. *Obstetrics & Gynecology*. 2015; 126(3)[DOI](#)
10. Hamilton C.A., Miller A., Casablanca Y., Horowitz N.S., Rungruang B., Krivak T.C., Richard S.D., Rodriguez N., Birrer M.J., Backes F.J., Geller M.A., Quinn M., Goodheart M.J., Mutch D.G., Kavanagh J.J., Maxwell G.L., Bookman M.A.. Clinicopathologic characteristics associated with long-term survival in advanced epithelial ovarian cancer: an NRG Oncology/Gynecologic Oncology Group ancillary data study. *Gynecologic Oncology*. 2018; 148(2)[DOI](#)
11. Zorn Kristin K., Tian Chunqiao, McGuire William P., Hoskins William J., Markman Maurie, Muggia Franco M., Rose Peter G., Ozols Robert F., Spriggs David, Armstrong Deborah K.. The prognostic value of pretreatment CA 125 in patients with advanced ovarian carcinoma. *Cancer*. 2009; 115(5)[DOI](#)
12. Schnack Tine H., Høgdall Estrid, Nedergaard Lotte, Høgdall Claus. Demographic Clinical and Prognostic Factors of Primary Ovarian Adenocarcinomas of Serous and Clear Cell Histology—A Comparative Study. *International Journal of Gynecological Cancer*. 2016; 26(1)[DOI](#)
13. Chan John K., Teoh Deanna, Hu Jessica M., Shin Jacob Y., Osann Kathryn, Kapp Daniel S.. Do clear cell ovarian carcinomas have poorer prognosis compared to other epithelial cell types? A study of 1411 clear cell ovarian cancers. *Gynecologic Oncology*. 2008; 109(3)[DOI](#)
14. Ye Shuang, Yang Jiaxin, You Yan, Cao Dongyan, Huang Huifang, Wu Ming, Chen Jie, Lang Jinghe, Shen Keng. Comparison of Clinical Characteristic and Prognosis between Ovarian Clear Cell Carcinoma and Serous Carcinoma: A 10-Year Cohort Study of Chinese Patients. *PLOS ONE*. 2015; 10(7)[DOI](#)
15. Melamed Alexander, Fink Günther, Wright Alexi A, Keating Nancy L, Gockley Allison A, del Carmen Marcela G, Schorge John O, Rauh-Hain J Alejandro. Effect of adoption of neoadjuvant chemotherapy for advanced ovarian cancer on all cause mortality: quasi-experimental study. *BMJ*. 2018. [DOI](#)
16. Rosen Barry, Laframboise Stephane, Ferguson Sarah, Dodge Jason, Bernardini Marcus, Murphy Joan, Segev Yakir, Sun Ping, Narod Steven A.. The impacts of neoadjuvant chemotherapy and of debulking surgery on survival from advanced ovarian cancer. *Gynecologic Oncology*. 2014; 134(3)[DOI](#)