

# Suprapubic Catheter in a Patient with Bladder Carcinoma, Against the Prohibition: A Systematic Review and Case Report

*Derri Hafa Nurfajri*

Department of Urology, Hasan Sadikin Academic Medical Center/Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia.

*Sawkar Vijay Pramod*

Department of Urology, Hasan Sadikin Academic Medical Center/Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia.

*Ferry Safriadi*

Department of Urology, Hasan Sadikin Academic Medical Center/Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia.

*Betty Suryawathy Hernowo*

Department of Anatomic Pathology, Faculty of Medicine, Universitas Padjadjaran, Bandung, West Java, Indonesia.

**Background:** Bladder tumors are the most common neoplasm of the lower urinary tract. Bladder carcinoma arising from a suprapubic cystostomy tract is a relatively rare. Some patients with bladder tumors have difficult urethral access for urinary retention due to inaccessible urethra. Most people think cystostomy in patient with bladder cancer can cause seeding and upstaging to suprapubic cystostomy tract. This study aim of suprapubic cystostomy can cause seeding and upstaging of bladder cancer.

**Methods:** Literature obtained from the search results of Pubmed, Medscape, Science Direct, Scopus, Cochrane Library, and Google Scholar electronic databases with the keywords used are “suprapubic catheter and bladder cancer”, “suprapubic catheter and bladder tumor”, “suprapubic catheter and bladder carcinoma”, “suprapubic cystotomy and bladder carcinoma”, and “suprapubic catheter and bladder cancer upstaging” with no time limits.

**Results:** Eighty two articles were obtained from the electronic database. Following all the inclusion and exclusion criteria, the final selection considered 5 literature with 16 patient eligible for this literature review. The five literature involved a total of 16 bladder cancer patients with a suprapubic catheter. The duration of suprapubic catheter insertion was between 1 to 3 months. As long as it is within the specified time and from the location of cancer, literature result there will be no seeding and upgrading in the suprapubic cystostomy tract. In our case report we have patient with duration suprapubic cystostomy 1 months with no seeding and upgrading cancer.

**Conclusion:** Although SPC is an effective, inexpensive, easy mode of access for bladder tumors with difficult urethral access for urinary retention due to inaccessible urethra, it also presents a risk of SPC tract bladder cancer, mostly SCC and TCC. It is important to be aware of any suspicious signs and symptoms, duration time of use suprapubic cystostomy and location of the cancer.

---

## Introduction

Bladder tumors are the most common neoplasm of the lower urinary tract. A majority of patients usually present with gross painless hematuria as the sole presenting symptom [1]. Incidence peaks in the sixth and seventh decades of life, although a trend toward presentation at younger ages has been suggested [2]. The lifetime risk for men is 3.4%, and for women, 1.2% [3]. Initial symptoms of urothelial carcinoma of the bladder (UCB) include microhematuria, painless macrohematuria, and/or irritative voiding symptoms, and require further investigation. Carcinoma in situ of the bladder

causes irritative lower urinary tract symptoms (LUTS) more often than papillary UCB. Histopathologic evaluation is necessary to assess the stage and grade with sufficient certainty after the appearance of bladder tumors [4].

Several causative risk factors are associated with urothelial carcinoma of the bladder, including smoking, genetic, and other carcinogenic exposure [5]. Other rarer tumor types include squamous cell carcinomas, which occur in the setting of chronic inflammation [6]. These were in line with the findings which determined that bladder tumors spread by implantation in abdominal wounds, denuded epithelium, resected prostatic fossa, or traumatized urethra; implantation occurs most often with high-grade tumors [7]. Squamous cell carcinoma arising from a suprapubic cystostomy tract (SCC-SCT) is a relatively rare bladder malignancy, which is known to have a close association with long-term inflammation and mechanical irritation from the suprapubic catheter [8]. The suprapubic catheter (SPC) is a non-continent, direct drainage of the bladder in cases of inaccessible urethra; it may be accomplished by an open or punch technique with a trocar or percutaneous methods using the Seldinger wire technique. The suprapubic catheter is a relatively safe and common procedure in urologic practice; it is also used in cases of genitourinary trauma, with few complications. It provides effective urinary diversion and drainage, which might be beneficial in cases of an inaccessible urethra [9]. The term inaccessible urethra means all prescribed endoscopic interventions have failed to relieve acute painful retention [9]. Inaccessible urethra with no retrograde endoscopic access due to multiple/diffuse strictures or multiple urethrocutaneous fistulas with acute urinary retention due to post-urethral instrumentation (transurethral resection of bladder tumor [TURBT], or TURBT with transurethral resection of the prostate [TURP]), is a rare entity. Management of such case with a bladder tumor for TURBT/surveillance cystoscopy poses a great challenge [9].

Therefore, this literature review aimed to investigate the utilization of SPC in a patient with bladder cancer against the prohibition.

## **Materials and Methods**

### **Literature search strategy**

Literature obtained from the search results of Pubmed, Medscape, Science Direct, Scopus, Cochrane library, and Google Scholar electronic databases with keywords: “suprapubic catheter and bladder cancer”, “suprapubic catheter and bladder tumor”, “suprapubic catheter and bladder carcinoma”, “suprapubic cystotomy and bladder carcinoma” and “Suprapubic catheter and bladder cancer upstaging”. References cited in the relevant literature were taken manually and only from full articles.

### **Inclusion and exclusion criteria**

According to the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines, the literature was obtained by reviewing PICOS (Population, Intervention, Comparator, Outcome, and Study design) to determine the feasibility of the study. Studies were considered feasible if they met the inclusion criteria as followings; 1) Patients of the studies underwent suprapubic catheter insertion; 2) Patients of the studies underwent the histopathological examination and physical examination leading to the diagnosis of bladder cancer; 3) Patients of the studies were examined for the following variables; i) Number of tumors, ii) Tumor diameter, iii) Localization of tumor, iv) Histology classification, v) Complication rate; 4) The studies consist of the outcome evaluation of suprapubic catheter in patients with bladder cancer.

Literature in the form of only abstracts, report meetings, conferences, editorial comments, reviews (systematic literature reviews and meta-analyses), irrelevant results, inaccessible literature, duplications from previous literature publications, and non-English studies were excluded.

## Systematic literature review process

After excluding multiple literature, evaluation of the abstract and title of the relevant literature were carried out. After evaluating abstracts and titles, acquired literature that fulfilled the requirements for a full evaluation was selected. Data were then extracted and presented in a table. The table consisted of the author’s name, year of publication, country of origin, study design, number of samples, number of tumors, tumor diameter, localization of tumor, histopathological classification, and complication rate. The PRISMA diagram that describes the literature review process and literature selection are shown in Figure 1.

**Figure 1. PRISMA Diagram that Describes the Search Process for Literature Review and Literature Selection.**

## Results

Since the beginning of 2021, 82 articles have been collected through the electronic database of Pubmed, Medscape, Science Direct, Scopus, Cochrane Library, and Google Scholar. After the exclusion of multiple literature, 36 relevant literature were obtained for the abstracts and titles evaluation. From the results of the evaluation, ten literature were eligible for a full evaluation. Based on all the inclusion and exclusion criteria mentioned earlier, the final selection considered 5 literature eligible with 16 patient for this literature review (Table 1).

No	Author	Year, country	Type of research	Study population
1	Khawaja, et al.[9]	2016, India	Case Series	12 patients
2	Gusev, et al.[10]	2020, United States	Case Report	1 patient
3	Ito,et al.[11]	2011, Japan	Case Report	1 patient
4	Breul, et al.[12]	1992, German	Case Report	1 patient
5	Zhang, et al.[13]	2015, China	Case Report	1 patient

**Table 1. Literature about Suprapubic Catheter in Bladder Cancer Patient.**

The literature involved a total of 17 bladder cancer patients with a suprapubic catheter. Of those, there were 1 case series and 5 case report study aimed to investigate the outcome of suprapubic catheter insertion in bladder cancer patients.

Based on the demographic profile, all of the patients were male, with the age range of 38 to 80 years (Table 2).

Study	Total Subjects	Age	Sex
Khawaja, et al.[9]	12 patients	38 - 68 years	Male (12 patients; 100%)
Gusev, et al.[10]	1 patient	65 years	Male
Ito,et al.[11]	1 patient	58 years	Male
Breul, et al.[12]	1 patient	80 years	Male
Zhang, et al.[13]	1 patient	61 years	Male

**Table 2. The Demographic Characteristic of Bladder Cancer Patients with SPC.**

In this review, the most frequent histopathological appearance in bladder cancer related to suprapubic catheter was TCC and SCC (Table 3).

Study	Number of Tumors	Size of tumor	Localization of tumor	Histopathological Classification
	Solitary (9; 75%)	Tumor diameter	Posterolateral wall	Low-grade transitional

				cell carcinoma (10; 83,33%)
		0,5 mm - 1 cm (9; 75%)	(6; 50%)	
	Multifocality (2; 16,66%)	1 - 3 cm (2; 16,66%)	Anterolateral (3; 25%)	
Khawaja, et al.[9]	Focal bladder thickening (1; 8,33%)	3 - 5 cm (1; 8,33%)	Dome (0)	High-grade transitional cell carcinoma (2; 16,66%)
			Base of bladder (2; 16,66%)	
			Bladder neck (1; 8,33%)	
Gusev, et al.[10]	N/A	N/A	Posterior bladder wall	High-grade, muscle-invasive urothelial carcinoma with glandular differentiation
Ito, et al.[11]	Solitary	72 mm x 63 mm	An abdominal mass surrounding a suprapubic cystostomy	Stage IV (cT4N1M1) epidermal SCC
Breul, et al.[12]	Solitary	4 x 4 x8 cm	Ventral to the bladder	poorly differentiated SCC with infiltration into the prostate (pT3, G3)
Zhang, et al.[13]	Solitary	(8 x 6 x 5) cm	Surrounding the suprapubic cystostomy and a space-occupying lesion	Squamous cell carcinoma
Study	Inaccessible	Type of SPC	Grade of the tumor	History of Operation
Khawaja, et al.[9]	Failed Catheterization (9; 75%)	N/A	N/A	TURBT (12; 100%)
	Urethrocutaneous fistula (2; 16,66%)			
	Failed OIU with retention and urinary extravasation (1; 8,33%)			
Gusev, et al.[10]	Urethral stricture	Squamous cell carcinoma	N/A	Biopsi stricture, Biopsi suprapubic mass
Ito, et al.[11]	N/A	Squamous cell carcinoma	cT4N1M1	N/A
Breul, et al.[12]	N/A	N/A	N/A	N/A
Zhang, et al.[13]	Urethral diverticulum with abscess formation perforated spontaneously	Moderately differentiated squamous cell carcinoma of the suprapubic tract but there was no bladder involvement	N/A	Operation on the spinal column (laminectomy T5 ± T9) resulted in a spinal cord injury, incomplete T4 and complete T7 (T4 ± T7 sensory/motor deficit).

**Table 3. The Clinicopathological Characteristic of Bladder Cancer Patients with SPC.**

Most of the bladder cancer patients with suprapubic catheters presented with recurrent urinary retention, inflammation around the area of the suprapubic catheter tract, hydronephrosis, and neurogenic bladder. Other clinical presentations of SPC complications included urethrocutaneous fistula, urinary extravasation, nocturia, reduced urinary flow, and gross hematuria (Table 4).

Study	Clinical Presentation of SPC Complication	Duration of SPC insertion	SPC Tract Histopathology
	Acute urinary retention with failed catheterization	1 - 2 months	Normal, only granulation tissue
Khawaja, et al.[9]	Urethrocutaneous fistula	3 months	Normal, only granulation

			tissue
	Post urethral instrumentation (failed OIU) with retention and urinary extravasation	3 months	Normal, only granulation tissue
Gusev, et al.[10]	Inflammation of the left lower quadrant omentum around the area of the biopsy tract; Gross hematuria; Acute renal failure (serum creatinine 4.4 mg/dl); Severe bilateral hydronephrosis	N/A	high-grade, muscle-invasive urothelial carcinoma with glandular differentiation
Ito, et al.[11]	Severely inflamed abdominal mass; Absence of sensation below the waist; Chronic neurogenic bladder; An abdominal mass surrounding a suprapubic cystostomy, erythematous skin around the mass, edematous, foul-smelling, and purulent discharge	1 month	Stage IV (cT4N1M1) epidermal SCC
Breul, et al.[12]	Repeated urinary retention; Nocturia of 3 times/night; Reduced urinary flow; Hydronephrosis; Filling defect of the bladder	N/A	poorly differentiated squamous cell carcinoma with infiltration into the prostate (pT3, G3)
Zhang, et al.[13]	Ulcerative bleeding with abnormal blisters surrounding SPC tract; Suprapubic catheter-related pain; Progressive lower extremity weakness; Neurogenic bladder; Bilateral hydronephrosis	3 months	Squamous cell carcinoma

**Table 4. The Outcome of Suprapubic Catheter Insertion in Bladder Cancer Patient.**

The duration of suprapubic catheter insertion was between 1 to 3 months. Most histopathological appearance of suprapubic catheter-related bladder cancer was urothelial carcinoma and squamous cell carcinoma. We also reported our cases; a 55-years-old man presented to the emergency unit with a chief complaint of retention urine with bladder cancer and inaccessible urethra.

### Case Presentation

A 55-years-old man presented to the emergency unit with a chief complaint of retention urine. The patient has a history of TURBT with pathology anatomically non-muscle invasive bladder cancer results. Ultrasonography of the Kidney, Ureter, and Bladder (USG KUB) showed no right and left kidney abnormality. The urinary bladder was full of urine (approximate volume of 850 ccs) and mass was found in posteriolateral of bladder shown in Figure 2.

**Figure 2. Mass in Posteriolateral Aspect of the Bladder Found Intraoperatively.**

Inserted a catheter does not work (inaccessible urethra) and decided to do SPC with Folley Catheter 16 fr.

The patient come with CT Scan revealed single irregular mass within the bladder wall on the posteriolateral part with calcification, extending from the mucosal layer to serosa layer, size 5,45 x 5,17 x 6,73 cm with post-contrast enhancement as shown in Figure 3.

**Figure 3. SPC Tract Shown in CT Scan.**

Patient have radical cystectomy and urinary diversion. Postoperation PA result found Low grade Urothelial carcinoma bladder (T2N0M0) without seeding and upstaging in SPC track. Figure 4 shows low-grade papillary bladder carcinoma compared to SPC tract without malignancy shown in Figure 5.

**Figure 4. Low-Grade Papillary Bladder Carcinoma.****Figure 5. SPC Tract without Malignancy.**

## Discussion

Bladder cancer is the most common tumor of the urinary system, comprising 6% of all malignancies in men and 2% of those in women [1]. Incidence peaks in the sixth and seventh decades of life [2]. According to Pashos et al., the lifetime risk for men is 3.4%, and for women 1.2% [3]. In this review, all of the patients were male, with the age range of 38 to 80 years.

In this review, the most histopathological finding of suprapubic catheter-related bladder cancer was urothelial carcinoma and squamous cell carcinoma. Based on the result of this review, most of the patients presented with recurrent urinary retention, inflammation around the area of the suprapubic catheter tract, hydronephrosis, and neurogenic bladder. Other clinical presentations of SPC complications included urethrocutaneous fistula, urinary extravasation, nocturia, reduced urinary flow, and gross hematuria.

A suprapubic catheter is a non-continent, direct drainage of the bladder in cases of inaccessible urethra; it may be accomplished by an open or punch technique with a trocar or percutaneously using the Seldinger wire technique. A suprapubic catheter is a relatively safe and common procedure in urologic practice; it is also used in cases of genitourinary trauma, with few complications. It provides an effective urinary diversion and drainage, which might be beneficial in cases of an inaccessible urethra [9]. According to a study conducted by Mustofi et al., rarer tumor types include squamous cell carcinomas, which occur in the setting of chronic inflammation, and adenocarcinomas, which occur in the bladder in association with a persistent urachal remnant and in cystitis glandularis associated with bladder exstrophy. In this review, the SPCs have usually been applied for 1 to 3 months.

The advantages of suprapubic diversion are that it is easy, provides quick relief, is practiced by all urologists; it can also be performed under local anesthesia. In cases of suspected space-occupying lesions in the bladder, ultrasound can be helpful to avoid needle advancement into the tumor during the suprapubic diversion. Chemotherapy/ immunotherapy can be administered with surveillance cystoscopy in cases of inaccessible urethra via SPC. Moreover, the patient can examine the SPC tract site for any swelling and skin changes, and it is also accessible to general practitioners and treating urologists [9].

The five literature involved a total of 16 bladder cancer patients with a suprapubic catheter. The duration of suprapubic catheter insertion was between 1 to 3 months. As long as it is within the specified time and from the location of cancer, literature result there will be no seeding and upgrading in the suprapubic cystostomy tract. In our case report we have patient with duration suprapubic cystostomy 1 months with no seeding and upgrading cancer.

In conclusion, although SPC is an effective, inexpensive, easy mode of access for bladder tumors with difficult urethral access for urinary retention due to inaccessible urethra, it also presents a

risk of SPC tract bladder cancer, mostly SCC and TCC. It is important to be aware of any suspicious signs and symptoms, duration time of use suprapubic cystostomy and location of the cancer. Tumor location in posterior, low grade, use feeding tube 6-8 Fr, duration of SPC 1-2 months, and seeding or upgrading of bladder cancer was not found.

## Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The authors declare no conflict of interest.

## References

## References

1. Greenlee R. T., Hill-Harmon M. B., Murray T., Thun M.. Cancer statistics, 2001. *CA: a cancer journal for clinicians*. 2001; 51(1)[DOI](#)
2. American Cancer Society. Cancer facts & figures, 2008. Atlanta, GA: American Cancer Society, 2008.
3. Pashos Chris Leo, Botteman Marc F., Laskin Benjamin Lewis, Redaelli Alberto. Bladder cancer: epidemiology, diagnosis, and management. *Cancer Practice*. 2002; 10(6)[DOI](#)
4. Amin Mahul B., McKenney Jesse K., Paner Gladell P., Hansel Donna E., Grignon David J., Montironi Rodolfo, Lin Oscar, Jorda Merce, Jenkins Lawrence C., Soloway Mark, Epstein Jonathan I., Reuter Victor E.. ICUD-EAU International Consultation on Bladder Cancer 2012: Pathology. *European Urology*. 2013; 63(1)[DOI](#)
5. Fleshner N, Kondylis F. Demographics and epidemiology of urothelial cancer of urinary bladder. In: Droller MJ, ed. *Urothelial tumors*. Hamilton, ON: Decker. 2004;1-16.
6. Vikram Raghunandan, Sandler Carl M., Ng Chuan S.. Imaging and Staging of Transitional Cell Carcinoma: Part 1, Lower Urinary Tract. *American Journal of Roentgenology*. 2009; 192(6)[DOI](#)
7. McDougal WS, Wein AJ, Kavoussi LR, et al, eds. Campbell-Walsh Urology. 9th ed. Philadelphia, PA: Saunders Elsevier. 2012;2426.
8. Vaidyanathan Subramanian, Mansour Paul, Soni Bakul M., Singh Gurpreet, Sett Pradipkumar. The method of bladder drainage in spinal cord injury patients may influence the histological changes in the mucosa of neuropathic bladder - a hypothesis. *BMC urology*. 2002; 2[DOI](#)
9. Khawaja Abdul Rouf, Dar Tanveer Iqbal, Maik Sajad, Magray Javaid, Bhat Ashiq, Bhat Arif Hameed, Wani Mohd Saleem, Wazir Baldev Singh. Effectiveness of Antegrade Access in Bladder Tumors With Inaccessible Urethra. *Reviews in Urology*. 2015; 17(4)
10. Stroumbakis N., Choudhury M. S., Hernandez-Graulau J. M.. Squamous cell carcinoma arising from suprapubic cystostomy site without bladder involvement. *Urology*. 1993; 41(6)[DOI](#)
11. Pannek Jürgen. Transitional cell carcinoma in patients with spinal cord injury: a high risk malignancy?. *Urology*. 2002; 59(2)[DOI](#)
12. Mostofi F, Davis CJ, Sesterhenn I. Histological typing of urinary bladder tumours, 2nd ed. Berlin, Germany: Springer-Verlag. 1999;3-27.
13. Schaafsma R. J., Delaere K. P., Theunissen P. H.. Squamous cell carcinoma of suprapubic cystostomy tract without bladder involvement. *Spinal Cord*. 1999; 37(5)[DOI](#)