

Surgical Treatment of Congenital Bronchiectasis in Childhood

Kh.X. Sultanov
M.M. Aliyev
I.R. Aytmuratov

Department of Pediatric Surgery, Uzbekistan.
Karakalpakistan Medical Institute, Uzbekistan.
Tashkent Pediatric Medical Institute, Uzbekistan.

Background and purpose: Congenital bronchiectasis (CB) is still an important problem for certain serious pulmonary infections. Surgical treatment of childhood CB has not been discussed extensively because of decline in prevalence and increased experience dealing with this disease. Although the incidence has declined over the past years in societies with high socioeconomic status, CB is still an important health problem in our country. This disease can be diffuse or focal. Diffuse CB is thought to occur owing to underlying systemic illness that causes abnormalities in airway clearance. For patients with bilateral disease, lung transplantation would be the only probable surgical approach. For patients with focal disease, conversely, the removal of the diseased lung had played the main role in the treatment of this disease. Recently, it has been shown that patients with an earlier and more localized disease can be successfully controlled by conservative approaches. It remains controversial as to which children would benefit from surgery and surgical points that may affect the outcome. Therefore, a retrospective study was conducted to evaluate the results of surgical treatment of CB in children.

Patients and Methods: We retrospectively reviewed the medical records of 79 consecutive children who underwent surgery for CB in our clinic between 2006 and 2021. Patient demographics, clinical features, radiologic examinations, details of surgery including type of resection, operative morbidity, mortality and outcomes were analysed.

Results: Seventy-nine patients underwent 83 pulmonary resections during the study period. The mean ages at diagnosis of CB and at the time of surgery were 8.10 ± 4.10 years (range, 1 to 17 years) and 9.70 ± 4.20 years (range, 2 to 18 years), respectively. There were 44 males (55.7%) and 35 females (44.3%). The causes of bronchiectasis were congenital hypoplasia of bronchi ($n = 31$), secondary bronchiectasis was additional nonspecific pulmonary infection ($n = 42$) and foreign body aspiration ($n = 6$). Chest X-rays, bronchography and bronchoscopy ($n = 79$), chest HRCT ($n = 61$), angiopulmonography ($n = 23$) were used, and pulmonary function tests ($n = 48$) were performed. The types of resections were segmentectomy (26.5%), lobectomy (35.4%), and lobectomy with segmentectomy (17.8%), bilobectomy (8.9%) and pneumonectomy (11.4%). Four patients with bilateral bronchiectasis subsequently required second operation for the other lung. Two of those had undergone right lower lobectomy, by one child underwent right upper lobe, and another left lower lobectomies. One patient with incomplete resection subsequently required second operation for ongoing bronchiectasis underwent complementary right pneumonectomy. Morphological study ($n=39$) Postoperative complications were encountered in 6 patients (7.6%). In terms of long-term outcomes of treatment 78 (98.7%) patients were followed-up at least 6 months up to 14 years. The postoperative status of the patients as follows: "well" in - 49 patients (60.7%), "improved" in 23 patients (29.5%), "orsened" in 6 patients (7.7%). And unfortunately two patients "died" (2.6%).

Conclusions: The decision for bronchiectasis surgery should be made in cooperation with the chest diseases unit. Anatomic localization of the disease should be mapped clearly by radiologic and angiographic investigations. A radiologic and morphologic evidence of reversal of airway abnormality has been shown in cases of post-infectious bronchiectasis and congenital defects of bronchial wall. The morbidity and mortality rates of bronchiectasis surgery are within acceptable ranges. Most of the children benefit from surgery, especially when complete excision is accomplished. Segmentectomy and lobectomy are well tolerated in

children without increase in morbidity and mortality. Therefore, resection of damaged part of the lung tissue may be preferred instead of removing high volume lung tissue.

Introduction

Congenital bronchiectasis (CB) is rather uncommon, and knowledge of this disorder is not very extensive. CB is defined as permanent and irreversible abnormal dilatation of bronchi, and is usually considered to be acquired. There is a type of bronchiectasis which is probably congenital in origin, although in most cases this is difficult to prove. Congenital cystic bronchiectasis is thought to be the result of developmental arrest. This may result in cyst formation, retention of fluid or air, and the risk of subsequent infection [1]. Congenital cystic bronchiectasis has been found almost exclusively in infants and young children [2, 3] and also has been reported accompanying other congenital anomalies [4], or in patients with deficiency of bronchial cartilages. Surgery was considered the only alternative method for the treatment of bronchiectasis in the early and mid years of this century. However, the indications for surgery have been limited in recent decades because of expectant results of conservative treatment. Although there are no prospective randomized trials comparing the efficacy of conservative and surgical treatment options, the indications of surgery for bronchiectasis can be considered in 2 main categories among children. Surgery can be performed for eradication in children suitable for total resection who do not respond to conservative treatment or for controlling the symptoms by removing the worst involved part in children with diffuse bronchiectasis. A retrospective series analysis was conducted to evaluate the results of surgical treatment according to the reasons and procedures in children with bronchiectasis.

Materials and Methods

We reviewed the medical records of all children who underwent surgery for bronchiectasis between 2006 and 2021 at Pediatric Surgery Department of the TashPMI and KMI. Medical records of 79 patients were analyzed for age, gender, clinical characteristics, radiologic findings and details of surgery including type of resection, operative morbidity, mortality, and overall outcomes.

Surgical treatment was considered if the symptoms persisted in spite of repeated courses of medical treatment including antibiotics, mucolytics, chest physiotherapy, bronchodilators, or steroids. Candidates were evaluated for the resectability of all diseased lung tissue or the expected benefit after excision of most affected parts of lungs with extensive disease.

Preoperative bronchography or chest high-resolution computed tomography (HRCT) was performed to evaluate the type, severity, and distribution of CB. Pulmonary function tests were done to estimate the lung function. Bronchoscopy was performed in all patients to rule out bronchial obstruction or stenosis caused by foreign body or endobronchial tumor.

Preoperative bronchoscopy was performed to clear bronchi and to prevent spillage of secretions during operation. Surgical access was obtained via posterolateral thoracotomy in all patients. The fourth or fifth intercostal spaces were used to reach the diseased lung area. The most common surgical procedures were formal segmentectomy and lobectomy, however, in some of the cases, when required lobectomy plus segmentectomy, bilobectomy and pneumonectomy was performed. A chest tube was placed and connected to an underwater seal system by Byulau in all patients. Postoperative management included intensive chest physiotherapy, analgesic, and antibiotics.

Complications were analyzed for intraoperative and postoperative periods. The outcome of surgery was evaluated at last follow-up visit and rated according to the degree of symptoms. Symptoms were categorized as follows: "well" when the patients are free of symptoms, "improved" when patient's symptoms reduced, "worse" if the patient's symptoms persist.

Results

Seventy-nine patients underwent 83 pulmonary resection operations with the diagnosis of bronchiectasis during the study period. The mean ages at diagnosis of bronchiectasis and at the time of surgery were 8.10 ± 4.10 years (range, 1 to 17 years) and 9.70 ± 4.20 years (range, 2 to 18 years), respectively. There were 44 males (55.7%) and 35 females (44.3%). The causes of bronchiectasis were nonspecific pulmonary infection (n = 42), congenital hypoplasia of bronchi (n = 31), and foreign body aspiration (n = 6). The details of comorbidities are shown in Table 1.

Comorbidities	No.	%
Oligophrenia	3	3.8
Congenital atrophy of the optic nerve	1	1.2
Chronic diseases of ENT organs	21	26.6
Chest wall deformity	5	6.3
Congenital heart disease	9	11.4
Aplasia or hypoplasia of the kidney	4	5
Hernias (diaphragmatic, inguinal, umbilical)	8	10.1
Congenital dislocation of the thigh	1	1.2
Total	52	54.8

Table 1. The Comorbidities with Bronchiectasis. Number of reported associated hereditary and inborn diseases in patients with bronchiectasis (n = 31).

The majority of comorbidities were diagnosed in the younger age group. Hidden leaking anomalies were identified with additional examination. It should be noted that in 11 (13.9%) children these diseases are detected with various combinations that significantly affect the type of treatment.

Initial radiologic evaluation was made through chest X-rays in all patients, and the diagnosis of bronchiectasis was confirmed by bronchography (n = 79) or chest HRCT (n = 61). Bronchography had been used in the early years of the study period. High-resolution CT has been preferred instead of bronchography afterwards. Pulmonary vascular angiography was performed in 23 patients to evaluate the functional and anatomical status of pulmonary circulation. Pulmonary function tests were performed in 48 children and showed pathologic results in 31.

The operative resections consisted of segmenectomy (26.5%), lobectomy (35.4%), and lobectomy with segmentectomy (17.8%), bilobectomy (8.9%) and pneumonectomy (11.4%). The details of surgical resection were given in Table 2.

Surgical Procedure		No. (%)
Segmentectomy		21 (26.5)
Lobectomy	Upper	7
	Middle	2
	Lower	19
Lobectomy+Segmentectomy	Left lower+	14 (17.8)
	IV-V segments	
Bilobectomy	Upper	2
	Lower	5
Pneumonectomy		9 (11.4)
Total		79 (100)

Table 2. Type of Surgery and Distribution of Resected Lung Parts.

A radiologic and morphologic evidence of reversal of airway abnormality has been shown in cases

of post- infectious bronchiectasis and congenital defects of the bronchial wall. All specimens underwent the pathologic examination that confirmed the diagnosis. Histopathological examination of all specimens showed varying degrees of bronchiectatic changes, including destruction of the lung parenchyma, congenital defects of segmental bronchus and bronchioles.

Four patients with bilateral bronchiectasis subsequently required second operation of the other lung. Two patients of them had undergone right lower lobectomy, by one childhoods underwent right upper lobe, another left lower lobectomies. One patient with incomplete resection subsequently required second operation for ongoing bronchiectasis underwent complementary right pneumonectomy.

Postoperative complication was encountered in 6 (7.6%) patients. One patient underwent reoperation for decortication in the second postoperative week. The remaining five children developed post-operative pneumonia, atelectasis and respiratory failure; those were managed by conservative means with bronchoscopy. Long-term results of treatment in terms of 6 months up to 14 years studied in 78 (98.7%) teens. The patients who have undergone surgery were postoperatively rated as follows: well, 48 patients (60.7%); improved 23 patients (29.5%); unchanged or worsened 6 patients (7.7%). Two patients died (2.6%); one of them in the early postoperative period, second was in the closest after the operating period with bilateral bronchiectasis. The details of the type of surgery are given in Table 3.

Procedure	Number of patients (%)	Results			
		Well	Improved	Worse	Exitus
Segmentectomy	21 (26.5)	17	4	-	-
Lobectomy	28 (35.4)	22	6	-	-
Lobectomy + Segmentectomy	14 (17.8)	7	5	1	1
Bilobectomy	7 (8.9)	2	3	1	1
Pneumonectomy	9 (11.4)	1	5	3	-
Total (%)	79 (100)	48 (60.7)	23 (29.5)	6 (7.6)	2 (2.6)

Table 3. Long-term results of Bronchiectasis Surgery According to Type of Surgical Resection.

Most of the patients who were well or clinically improved had completely resectable lesions. Segmentectomy and lobectomy are well tolerated in children without increase in morbidity and mortality. Therefore, resection of damaged part of the lung tissue may be preferred instead of removing much volume lung tissue ($P < 0.05$).

Discussion

Bronchiectasis was uniformly fatal when it was described in detail by Laennec in 1819. Demonstration of abnormalities of bronchial architecture by using nebulized bismuth powder and iodized oil allowed further understanding of the pathogenesis of bronchiectasis. In modern description, bronchiectasis is characterized by abnormal, irreversible dilatation of the bronchi in association with a variable degree of chronic bronchitis, pneumonitis, or other pathologic changes.

The common cause of bronchiectasis is bacterial infection usually after pneumonia. Other causes include pertussis, viral infections, and obstruction secondary to foreign body aspiration, tracheobronchial aspiration, congenital lesions and immune deficiency. Cystic fibrosis is the most common cause of bronchiectasis especially in the white population of Europe and North America. Nonspecific pulmonary infection was the most frequent cause of bronchiectasis in our series. The prevalence of bronchiectasis decreased significantly over recent decades. In the base of some of bronchiectasis lays the hypoplasia of the bronchial wall elements, bronchioles and alveolus in whole

or limited area of the lung. Antimicrobial therapy and immunization against viral and bacterial agents significantly decreased the incidence of bronchiectasis. Early recognition of foreign bodies with bronchoscopy also decreased the incidence of postobstructive bronchiectasis. However, the incidence of bronchiectasis is still high in developing countries.

The initial treatment of bronchiectasis is conservative and includes reducing airway obstruction and elimination of microorganisms from the lower respiratory tract. Therefore, medical therapy for bronchiectasis consists of antimicrobial therapy, postural physiotherapy, bronchodilators, corticosteroids and bronchoscopic cleaning of the tracheobronchial tree (TBT).

The main rationale for surgical treatment of bronchiectasis is to increase the quality of children's lives and protect them from complications such as empyema, hemoptysis, and lung abscess. The successful results can be achieved only by careful and proper selection of candidates for surgery. Instead of following rigid criteria for patient selection, each child should be evaluated separately for surgery.

The indications for surgery are not fully established in children. However, there seems to be agreement that the appropriate antibiotic therapy and chest physiotherapy should be tried for a period. In adults, indications for surgery have been reported to be resistance to appropriate antibiotic therapy in addition to an extensive postural drainage and chest physiotherapy program for a reasonable period of at least 2 years. In children, in addition to the above-mentioned criteria, growth retardation, failure to thrive, inability to follow the educational program because of frequent infections, and socioeconomic status of the family should also be considered when deciding on surgical treatment.

Since all diseased parts of the lung can be resected, the decision for surgery is relatively easy in children with localized bronchiectasis. The presence of accompanying complications such as growth retardation, abscess, and hemoptysis help surgeons in their decision for operation. However, when the bronchiectasis diffusely affects multiple lobes or segments and sometimes both lungs, the operative decision is quite difficult. Especially in the presence of comorbidities such as deformation of thorax and congenital heart diseases, total resection of all affected tissue usually is impossible. All candidates of surgery should be free of an ongoing pulmonary infection at the time of operation. Broad-spectrum antibiotics should be given in the preoperative period. Bronchoscopy should be performed in all candidates to exclude bronchial obstruction caused by a foreign body. Bronchoscopy should be also done just before the operation to clear airways. In our series, a rigid or flexible bronchoscopy, or both, was performed for the removal of secretions and the detection of a foreign body or an endobronchial lesion.

The localization of disease should be evaluated radiologically. Bronchography had been used extensively for both diagnostic and follow-up purposes in bronchiectasis. Although it was very sensitive and informative, it has been almost totally abandoned today in children. High-resolution chest CT has replaced the bronchography. Angiography has been recommended to find out whether the diseased area is perfused and detected of pulmonary vessels hypoplasia. It has been suggested that only nonperfusing lung and pulmonary vessels hypoplasia areas should be resected. Pulmonary function tests may not be required in children with localized bronchiectasis. However, it should be performed in cases of extensive involvement or in patients undergoing reoperations.

The aim at operation is to excise all diseased lung areas whenever possible and to preserve as much healthy lung parenchyma as possible. It is known that even only 2 or 3 preserved segments can fill the hemithorax, if operation had performed till 8 age old. All resections should be performed within anatomic limits. Lobectomy was the most frequently performed operation in our series. Left lower lobe was the frequently affected area as seen usually in bronchiectasis. Pneumonectomy was the second most frequent operation. Although pneumonectomy has been accepted to result more complications and mortality than that of other limited resections, complications has occurred after removing much volume lung tissue, such as lobectomy plus segmentectomy, bilobectomy and

pneumonectomy in long-term period, and it has been well tolerated in children in the current series.

The success of surgery for CB is directly related to the extension of the disease, underlying cause of bronchiectasis, and the completeness of the resection. Most of the children benefited from bronchiectasis surgery with regression of symptoms. Bilateral bronchiectasis exposed a high risk of mortality with all deaths occurring in these children. We could not find the exact cause of deaths in this patient, because no autopsy was allowed. Patients with bilaterally bronchiectasis should undergo a complete cardiologic evaluation for possible accompanying intrinsic abnormalities before operation.

Total excision of the diseased lung areas was accomplished in most children resulting in an apparent improvement in symptoms. However, symptoms have been persistent in children who had incomplete resection. Whether in the same or other site, the presence of residual diseased lung area has the potential to produce symptoms.

The decision for surgical intervention should be made in cooperation with the chest diseases unit. Anatomic localization of the disease should be mapped clearly by radiologic investigations. In addition, radiologic and morphologic evidence of reversal of airway abnormality has been shown in cases of post-infectious bronchiectasis. The morbidity and mortality rates of bronchiectasis surgery are within acceptable ranges. Most of the children benefit from the surgery, especially when total excision of the all diseased lung areas can be accomplished. Segmentectomy and lobectomy are well tolerated in children without increase in morbidity and mortality. Therefore, resection of damaged part of the lung tissue may be preferred instead of removing much volume lung tissue, but pneumonectomy may be preferred instead of leaving residual disease in the absence of contralateral lung involvement.

References

References

1. Alshamiri KM, Abbod HB. Congenital cystic adenomatoid malformation. *International Journal of Pediatrics & Adolescent Medicine*. 2017; 4(4)[DOI](#)
2. Delgado-Peña YP, Torrent-Vernetta A, Sacoto G, Mir-Messa I, Rovira-Amigo S, Gartner S, Moreno-Galdó A, Molino-Gahete JA, Castillo-Salinas F. Pulmonary hypoplasia: An analysis of cases over a 20-year period. *Anales De Pediatría (Barcelona, Spain: 2003)*. 2016; 85(2)[DOI](#)
3. Brower KS, Del Vecchio MT, Aronoff SC. The etiologies of non-CF bronchiectasis in childhood: a systematic review of 989 subjects. *BMC Pediatrics*. 2014; 14(1)[DOI](#)
4. DeBoer EM, Swiercz W, Heltshe SL, Anthony MM, Szeffler P, Klein R, Strain J, Brody AS, Sagel SD. Automated CT Scan Scores of Bronchiectasis and Air Trapping in Cystic Fibrosis. *Chest*. 2014; 145(3)[DOI](#)