



Awareness of Oral Cancer and OPMDs among Patients Attending the University Dental Hospital, Peradeniya, Sri Lanka

E. A. Shanika D. Somathunga

D. M. Sithara H. Dissanayaka

D. R. Dilini L. Ratnayake

Ruwan D. Jayasinghe

Dept. of Oral Medicine and Periodontology, Faculty of Dental Sciences, University of Peradeniya, Sri Lanka

Objective: To assess awareness of oral cancer and OPMDs among patients attending for dental treatment at the University Dental Hospital, Peradeniya.

Methods and materials: A self-administered questionnaire was used to collect information from 500 randomly selected outpatients attended the University Dental Hospital, Peradeniya, Sri Lanka. The questionnaire included questions to ascertain information on socio-demographic parameters, awareness of oral cancer and OPMDs, habits of betel chewing, smoking and alcohol consumption.

Results: 43.2% of the respondents were satisfactorily aware of the possibility of occurrence of oral cancer and OPMDs. From all the individuals who were participated to this questionnaire, 16 (3.2%) of individuals have a habit of betel chewing, 4.2% of individuals have a habit of smoking and 8.3% of individuals have a habit of alcohol consumption, respectively while having a satisfactory awareness about oral cancer and OPMDs.

Conclusions: This survey revealed that most of the patients attending the hospital were not in a satisfied level of awareness about oral cancer and OPMDs. Knowledge about the causal relationships with betel chewing was low compared to tobacco smoking and alcohol consumption.

Introduction

Oral cancer (OCA) poses a huge challenge to the wellbeing of people worldwide, particularly across South Asia. Oral cancer (OCA) is a malignant neoplasm of lip, tongue and mouth. Taken together, these cancers were estimated to affect 354,864 people worldwide in 2018, with 177,384 deaths [1]. Fifty-six per cent of the world's oral and pharyngeal cancer burden (excluding nasopharyngeal carcinoma which has different aetiology and biology) is from Asia where overall incidence of oral cancer for men was estimated at 14.8 per 100,000 population per annum in 2018 [1]. Oral cancer is the most common cancer among men in Sri Lanka, eighth amongst women, and second overall [2-3]. Oral cancer accounts for nearly 10% of reported cancers with the highest mortality rates of all cancers in Sri Lanka [3]. Yet oral cancer is entirely preventable.

Oral cancer is usually preceded with an oral potentially malignant disorder (OPMD). An OPMD is defined as a clinical disorder, usually characterized by visible alterations in the oral mucosa, which

may be a generalized change or appear as a more or less circumscribed lesion or lesions, in which there is increased risk for an oral squamous cell carcinoma to develop. Such carcinomas may develop within or without any discrete lesion/s [4]. Leukoplakia, OSF and lichen planus are the most common OPMDs in Sri Lanka. Erythroplakia is rare but has a very high malignant transformation rate. The transformation rate of leukoplakia is very variable, depending on the definitions used, the population studied, study design and length of follow up [5-6-7-8]. Global prevalence of OPMD is estimated to range between 1 and 5% [9-10]. Mello et al [10] estimated the prevalence of OPMDs to be 10.54% for Asian populations but these findings vary by type of OPMD: The prevalence of oral submucous fibrosis (OSMF) is estimated to be 4.96%, Oral Leukoplakia (OL) 4.11%, and 0.17% prevalence for erythroplakia [10]. In Sri Lanka, the prevalence of OL and of OSF was reported as 2% and 1.5% respectively among the 35-44 and 65+ age groups [11]. In a high risk population of tea estate labourers of Sri Lanka, the prevalence for OPMD was reported as 6.7% with 4.61% prevalence for OL and 1.64% for OSF [12]. Such wide geographical variations are due to life styles specific to the country or the region. Betel quid chewing with or without tobacco, chewing of areca nut, either traditional or commercially prepared forms are major contributory factors for oral cancer and OPMD in the South Asian region.

As management of oral cancer in Sri Lanka is costly

[13] and seriously affects quality of life primary prevention and early detection of OPMDs is essential if we are to reduce the incidence and to reduce the impact of this dreadful disease on population health and healthcare costs. The stage of diagnosis is important for survival. Oral cancers have a high likelihood of cure when detected at an early stage [14]. The lack of awareness on oral cancer and oral potentially malignant disorders (OPMD) and their risk factors among public as well as among health care professionals [15] is an important reason for late detection of these conditions [16].

Previous studies have demonstrated that the knowledge on oral cancer among people's better compared to OPMD [17-18-19]. Their knowledge on risk factors varies. In some studies there is an overall lack of awareness on risk factors [19] whereas some other have shown a low awareness on smoking and alcohol compared to betel chewing [17-18].

Education programs aiming at minimizing the oral cancer risk has to be developed using the existing knowledge and the gaps in the knowledge. Multiple educational programs are conducted in Sri Lanka by different organizations. Therefore, the primary aim of this study was to assess the awareness and knowledge of oral cancers and oral potentially malignant disorders and their risk factors among the patients attending the university dental hospital, Peradeniya, Sri Lanka.

Materials and Methods

A hospital based cross sectional study was conducted at the Out Patients Department (OPD) of the University Dental Hospital, Peradeniya, Sri Lanka. Randomly selected 500 OPD attendees were recruited to the study.

Self-administered pre tested questionnaire was given to the study participants while they were waiting for treatment. The self-administered questionnaire includes questions to ascertain information on socio-demographic of betel chewing, smoking and alcohol consumption.

Information sheet with necessary details was provided to the participants and informed consent was obtained from all participants prior to the administration of the questionnaire. Ethics approval was obtained from the Ethics Review Committee of the Faculty of Dental Sciences, University of Peradeniya, Sri Lanka. Data were analyzed using SPSS version 21 and chi-squared test was applied to compare groups.

Results

The study population consisted of 275 females and 225 males with an age range of 14-86 years and an average age of 36 (Table 1). Most of the participants had completed their education up to secondary education (71%) whereas only 3% had education up to primary education. Socio-demographic data of the study participants are summarized in the Table 1.

	Number	Percentage (%)
Gender		
Male	225	45
Female	275	55
Age		
14-24	139	27.8
25-34	95	19
35-44	93	18.6
45-54	91	18.2
55-64	45	9
Above 64	37	7.4
Ethnicity		
Major ethnic group	428	85.6
Minor ethnic groups	72	14.4

Table 1: Distribution of Respondents According to the Socio-demographic Data.

Level of awareness was considered satisfactory when an individual obtained a score of 50% or more. Accordingly, forty six percent of the respondents had satisfactory level of awareness and the level of awareness was significantly associated with the level of education ($\chi^2 = 14.296$, $p = 0.001$) of the study participants. Binomial logistic regression was performed to ascertain the adjusted effects of age, gender, ethnicity, occupation and level of education on the likelihood that study participants possessed a satisfactory level of awareness. The logistic regression model was statistically significant ($\chi^2 = 29.46$, $P = 0.003$). However the model explained only 7.6% (Nagelkerke R^2) of the variance in awareness level and correctly classified 60.4% of cases who possessed a satisfactory and non-satisfactory level of awareness. According to the model those who have obtained their education up to G.C.E. O/L or G.C.E. A/L (secondary education) ($OR = 5.41$, 95% $CI = 1.46-19.98$, $p = 0.01$) and primary education possessed a higher level of awareness ($OR = 1.82$, 95% $CI = 1.14-2.28$, $p = 0.012$) when compared with the individuals who have possessed their tertiary education. Most of them have obtained the awareness from TV/radio, newspapers, posters or banners whereas only around 10% have received it from healthcare workers (Table 2).

Sources of awareness about Oral Cancer and OPMD	Frequency	Percentage (%)
No response	171	34.2
From Family Members	8	1.6
From Friends	8	1.6
From a Physician	24	4.8
From a Dental Surgeon	29	5.8
From Midwife	9	1.8
By Newspaper	22	4.4
By T.V. /Radio	71	14.2
By Other ways (poster, banner)	22	4.4
Two or More responses	136	27.2
Total	500	100

Table 2: Sources of Awareness about Oral Cancer and OPMDs.

Some questions were directed to ascertain the awareness about the link between oral cancer and OPMDs with various habits including betel chewing 16 (3.2%), tobacco smoking 4.2% and alcohol consumption 8.3%. Out of 500 subjects 2.8% have used betel quid with all ingredients (betel leaf, areca nut, tobacco and lime) while 2.6% chewed without tobacco. Furthermore habit of betel chewing was significantly associated with the level of awareness on oral cancer and OPMD ($P=0.008$) (Table 3).

Major risk factors	X ²	df	P value
Betel Chewing	6.916	1	0.008
Smoking	0.824	1	0.364
Alcohol	1.447	1	0.229

Table 3: Association between Habits and Level of Awareness on Oral Cancer and OPMD.

Discussion

Ministry of Health of Sri Lanka, Faculty of Dental Sciences of the University of Peradeniya, Sri Lanka and many other governmental and non-governmental organizations and institutions conduct multiple programs using multiple methods of communication to educate the people of Sri Lanka on oral cancer prevention especially on the prevention of risk habits. Yet, oral cancer is a major public health problem in Sri Lanka [3]. Even though there is a reduction in smoking, number of people who use betel quid is increasing [20]. Oral cancer is usually preceded by an oral potentially malignant disorder (OPMD), especially in the south and south East Asia. Global prevalence of OPMD is estimated to be 1 -5% (Napier and Speight, 2008, Mello et al., 2018). Prevalence of OPMD is higher in South & South East Asia [21-22]. In Sri Lanka, the prevalence of oral leukoplakia (OL) and oral sub mucous fibrosis (OSF) was reported as 2% and 1.5% respectively among the 35-44 and 65+ age groups [11]. Even though oral cancer occurs in an anatomical site which is visible to both the affected individual and to an examining clinician, most patients with an OCA report late to the hospitals, with advanced disease. Every year around 2500 new cases of OCA are diagnosed in Sri Lanka where almost all patients go on to receive treatment from state sector hospitals. Sri Lanka is a country with 100% free health services. According to the National Cancer Incidence data year 2011, more than 76.5% of the total reported oral cancer cases were in UICC/AJCC stages 111 and 1V [3]. This is one of the most important poor prognostic marker of oral cavity cancers. Lack of awareness about OPMD and oral cancer is one reasons for late diagnosis. Therefore early identification of oral cavity cancer and OPMD by health care professionals and by people themselves will be a great advantage in the management of oral cavity cancer.

According to the findings of current survey, only 43.2% of participants had a satisfactory level of awareness on oral cancer and OPMDs. Knowledge on oral cancer was much better than that of the OPMD. Oral cancer/ OPMD awareness among people varies considerably in literature. Some have shown a poor knowledge like ours [18] whereas some other have shown a better knowledge [23-24]. A similar study conducted in the same hospital 15 years ago, awareness on oral cancer was 95% whereas for precancer it was 45% highlighting a significant reduction in the level of awareness among the patients attending to the Dental Hospital, Peradeniya [17]. Management of oral cavity cancer is not an easy task, most of the time leading to poor prognosis, poor quality of life of the patients and are costly whereas the management of OPMD is much easier, economical with better outcomes [13]. Timely diagnosis of OPMD and proper management can help in preventing malignant transformation. In this context it is important to note that more than 55% of the respondents were without a satisfactory level of knowledge on OPMDs and oral cancers and awareness on oral cancer/ OPMD has reduced over the years. This information is an eye opener to the health care professional especially public health specialists of this country and points to the dire need for educational programs to enlighten the public about this dreadful disease. Surprisingly we observed a lower level of awareness in people with higher level of education.

Even though the manufacturing and selling of smokeless tobacco is banned in Sri Lanka, still betel quid chewing with smokeless tobacco and areca nut is a very common habit. Many activities using different approaches and different groups have been tried in smokeless tobacco cessation [25] but it is difficult task due to its relationship with social, religious and cultural practices in Sri Lanka. Findings of current survey revealed that the knowledge about the causal relationship with betel chewing was low compared to tobacco smoking and alcohol consumption. But the previous hospital-based study conducted in Sri Lanka found, awareness of the causal relationship of tobacco in the betel quid and occurrence of oral cancer was high compared to tobacco smoking and alcohol consumption [17] demonstrating a declining awareness among people.

Mass media has been identified as an effective tool in educating the people regarding oral cancer and OPMD [26]. There had been many mass media programs organized by various institutions aiming public health education including oral cancer prevention in Sri Lanka. This survey showed that the mass media play a pivotal role in educating the public about oral cancer. Since oral cancer and OPMDs are commonly found in socially and economically deprived strata of the population, it is doubtful whether modes of mass media are readily accessible to the underprivileged groups. The public should be educated and positively encouraged to participate in educational programs on cancer prevention with the hope of minimizing the incidence of this dreadful disease in Sri Lanka. Only a small group of participants have received awareness through dental surgeons/medical officers which has been noted in some other studies as well [23-24]. Health care professionals have a duty and a responsibility in providing health education to their patients. Therefore health care professionals must take an active role in educating the people on oral cancer/ OPMDs.

Although the present study was a hospital-based survey in an urban area, findings of the study revealed some important aspects of the awareness of oral cancer and OPMDs and their risk factors. It demonstrate that the awareness is not adequate and more importantly awareness on OPMD/ Oral cancer as well as risk factors has declined over period of 15 years compared to a similar study. Findings of our study revealed that only a small percentage of participants had received their knowledge on OPMD/Oral Cancer from medical officers or dental surgeons. Major limitation of our study is that it was a hospital based study therefore subjects that we have included are biased and may have better awareness than the people who do not come in to hospitals. Most of the patients coming to this hospital are from urban areas. Most of our patients are either young or middle aged. Therefore it may not be appropriate to generalize the findings hence a well-designed population-based study with a larger sample is need to see the awareness in the population.

Recommendations

Within those limitations following recommendations can be made from the findings of this study. As the study show a relatively low awareness, it is important to improve the public awareness on OPMD/ Oral cancer and their risk factors. Existing awareness programs need to be reassessed and well-structured, socially and culturally acceptable educational programs should be designed and carried out in the society. Health care professionals should be educated and motivated to engage in patient education regarding the OPMD and oral cancer prevention.

References

References

1. Ferlay J, E. M, Lam F, Colombet M, Mery L, Piñeros M, Znaor A, Soerjomataram I, Bray F. Global Cancer Observatory: Cancer Today. year 2018. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today>.
2. National Cancer Control Programme, Ministry of Health, Nutrition & Indigenous Medicine, Sri Lanka. *Cancer Incidence Data: Sri Lanka Year 2011. Colombo: NCCP. 13th*

Publication.

3. National Cancer Control Programme, Ministry of Health, Nutrition & Indigenous Medicine, Sri Lanka - 2014, 2014. *Cancer Incidence Data Sri Lanka 2014*. [Www.Nccp.Health.Gov.Lk/Index.Php/Publications/Cancer-Registries](http://www.Nccp.Health.Gov.Lk/Index.Php/Publications/Cancer-Registries).
4. van der Waal Isaac. Potentially malignant disorders of the oral and oropharyngeal mucosa; terminology, classification and present concepts of management. *Oral Oncology*. 2009; 45(4-5)[DOI](#)
5. Warnakulasuriya S., Ariyawardana A.. Malignant transformation of oral leukoplakia: a systematic review of observational studies. *Journal of Oral Pathology & Medicine*. 2015; 45(3)[DOI](#)
6. Farah Camile S., Jessri Maryam, Bennett Nigel C., Dalley Andrew J., Shearston Kate D., Fox Simon A.. Exome sequencing of oral leukoplakia and oral squamous cell carcinoma implicates DNA damage repair gene defects in malignant transformation. *Oral Oncology*. 2019; 96[DOI](#)
7. Shearston Kate, Fateh Behrooz, Tai Shixiong, Hove Dzikamai, Farah Camile S.. Malignant transformation rate of oral leukoplakia in an Australian population. *Journal of Oral Pathology & Medicine*. 2019; 48(7)[DOI](#)
8. Shearston Kate, Fateh Behrooz, Tai Shixiong, Hove Dzikamai, Farah Camile S.. Oral lichenoid dysplasia and not oral lichen planus undergoes malignant transformation at high rates. *Journal of Oral Pathology & Medicine*. 2019; 48(7)[DOI](#)
9. Napier Séamus S., Speight Paul M.. Natural history of potentially malignant oral lesions and conditions: an overview of the literature. *Journal of Oral Pathology & Medicine*. 2007; 37(1)[DOI](#)
10. Mello Fernanda Weber, Miguel Andressa Fernanda Paza, Dutra Kamile Leonardi, Porporatti André Luís, Warnakulasuriya Saman, Guerra Eliete Neves Silva, Rivero Elena Riet Correa. Prevalence of oral potentially malignant disorders: A systematic review and meta-analysis. *Journal of Oral Pathology & Medicine*. 2018; 47(7)[DOI](#)
11. Ministry of Health Sri Lanka , -. 2019. National Oral Health Survey, Sri Lanka (2015/2016), Colombo: Ministry of Health, Nutrition and Indigenous Medicine (Sri Lanka); 2018. Available at <http://www.moh.gov.lk>.
12. Ariyawardana A., Sitheequ M. A. M., Ranasinghe A. W., Perera I., Tilakaratne W. M., Amaratunga E. A. P. D., Yang Yi-Hsin, Warnakulasuriya S.. Prevalence of oral cancer and pre-cancer and associated risk factors among tea estate workers in the central Sri Lanka. *Journal of Oral Pathology & Medicine*. 2007; 36(10)[DOI](#)
13. Amarasinghe H, Jayasinghe R. D, Dharmagunawardene D, Attygalla M, Scuffham P. A, Johnson N, Kularatna S. Economic burden of managing oral cancer patients in Sri Lanka: a cross-sectional hospital -based costing study. *BMJ Open*. 2019; 9:e027661.
14. Le Campion Anna Carolina Omena Vasconcellos, Ribeiro Camila Maria Beder, Luiz Ronir Raggio, da Silva Júnior Francisco Feliciano, Barros Herbert Charles Silva, dos Santos Karine de Cássia Batista, Ferreira Stefania Jeronimo, Gonçalves Lucio Souza, Ferreira Sonia Maria Soares. Low Survival Rates of Oral and Oropharyngeal Squamous Cell Carcinoma. *International Journal of Dentistry*. 2017; 2017[DOI](#)
15. Jayasinghe R D, Sherminie L P G, Amarasinghe H, Sitheequ M A. Level of awareness of oral cancer and oral potentially malignant disorders among medical and dental undergraduates. *Ceylon Medical Journal*. 2016; 61(2)[DOI](#)
16. Amarasinghe Hemantha K., Usgodaarachchi Udaya S., Johnson Newell W., Lalloo Ratilal, Warnakulasuriya Saman. Public awareness of oral cancer, of oral potentially malignant disorders and of their risk factors in some rural populations in Sri Lanka. *Community Dentistry and Oral Epidemiology*. 2010; 38(6)[DOI](#)
17. Ariyawardana A, Vithanaarachchi N. Awareness of Oral Cancer and Precancer among Patients Attending a Hospital in Sri Lanka. *Asian Pacific Journal of Cancer Prevention*. 2005; 6:58-61.
18. Formosa Joseph, Jenner Rachel, Nguyen-Thi My-Duyen, Stephens Caitlin, Wilson Corey, Ariyawardana Anura. Awareness and Knowledge of Oral Cancer and Potentially Malignant Oral Disorders among Dental Patients in Far North Queensland, Australia. *Asian Pacific*

- Journal of Cancer Prevention*. 2015; 16(10):4429-4434.
19. Shivakumar KM, Kadashetti Vidya, Choudhary Minal, Patil Swati, Gawnde Madhuri. Awareness and knowledge of tobacco associated risk of development of oral cancer and oral potentially malignant disorders among patients visiting a dental college. *Journal of Family Medicine and Primary Care*. 2020; 9(5)[DOI](#)
 20. www.searo.who.int/tobacco/documents.
 21. Chung Ching-Hung, Yang Yi-Hsin, Wang Tung-Yiu, Shieh Tien-Yu, Warnakulasuriya Saman. Oral precancerous disorders associated with areca quid chewing, smoking, and alcohol drinking in southern Taiwan. *Journal of Oral Pathology and Medicine*. 2005; 34(8)[DOI](#)
 22. Thomas Steven J., Harris Ross, Ness Andrew R., Tauro Joachim, MacLennan Robert, Howes Noah, Bain Christopher J.. Betel quid not containing tobacco and oral leukoplakia: A report on a cross-sectional study in Papua New Guinea and a meta-analysis of current evidence. *International Journal of Cancer*. 2008; 123(8)[DOI](#)
 23. Prado NS., Bonan RF., Leonel AC., Castro uF., Carvalho EJ., Silveira FM., Perez DE.. Awareness on oral cancer among patients attending dental school clinics in Brazil. *Medicina Oral Patología Oral y Cirugía Bucal*. 2020. [DOI](#)
 24. Villa Alessandro, Kreimer Aimee R., Pasi Massimo, Polimeni Antonella, Cicciù Domenico, Strohenger Laura, Gherlone Enrico, Abati Silvio. Oral Cancer Knowledge: A Survey Administered to Patients in Dental Departments at Large Italian Hospitals. *Journal of Cancer Education*. 2011; 26(3)[DOI](#)
 25. Hettiarachchi PVKS, Jayasooriya Primal, Amarasinghe Hemantha, Siriwardena BSMS, Wijerathne Deepashika, Kithalawa arachchi Samantha, Tilakaratne WM, Jayasinghe Ruwan. Knowledge and Attitudes of Nursing Students Towards Smokeless Tobacco and Areca Nut Control in Central Province of Sri Lanka. *Asian Pacific Journal of Cancer Care*. 2020; 5(3)[DOI](#)
 26. Shrestha Ashish, Rimal Jyotsna. Effectiveness of a Mass Media Campaign on Oral Carcinogens and Their Effects on the Oral Cavity. *Asian Pacific Journal of Cancer Prevention*. 2018; 19(3)[DOI](#)